



Galveston County POTENTIALLY PREVENTABLE HOSPITALIZATIONS

www.dshs.state.tx.us/ph

- From 2005-2008, adult residents (18+) of **Galveston County** received **\$459,688,908** in charges for hospitalizations that were potentially preventable.
- Hospitalizations for the conditions below are called “**potentially preventable,**” because **if the individual had access to and cooperated with appropriate outpatient healthcare, the hospitalization would likely not have occurred.**

Potentially Preventable Hospitalizations (2005-2008)				
Hospitalizations for Adult Residents of Galveston County	Number of Hospitalizations	Average Hospital Charge	Total Hospital Charges	Average \$ Impact for All Adult County Residents
Bacterial Pneumonia	3,550	\$28,357	\$100,668,244	\$460
Dehydration	791	\$17,026	\$13,467,701	\$62
Urinary Tract Infection	2,335	\$19,059	\$44,502,640	\$204
Angina (without procedures)	177	\$18,281	\$3,235,736	\$15
Congestive Heart Failure	4,590	\$30,040	\$137,885,381	\$631
Hypertension (High Blood Pressure)	734	\$18,738	\$13,753,444	\$63
Asthma	1,042	\$21,750	\$22,663,029	\$104
Chronic Obstructive Pulmonary Disease	2,285	\$25,493	\$58,250,704	\$266
Diabetes Short-term Complications	513	\$24,373	\$12,503,165	\$57
Diabetes Long-term Complications	1,561	\$33,798	\$52,758,864	\$241
TOTAL	17,578	\$26,151	\$459,688,908	\$2,102

Source: Center for Health Statistics, Texas Department of State Health Services

The purpose of this information is to assist in improving health care and reducing health care costs for Galveston County residents. This information is not an evaluation of hospitals or other health care providers.

Access to and cooperation with outpatient health care potentially prevents hospitalizations for the conditions below:

Bacterial Pneumonia is a serious inflammation of the lungs caused by an infection. Bacterial pneumonia primarily impacts older adults. **Communities can potentially prevent hospitalizations by encouraging older adults and other high risk individuals to get vaccinated for bacterial pneumonia and by encouraging the appropriate use of antibiotics.**

Dehydration means the body does not have enough fluid to function well. Dehydration primarily impacts older adults or institutionalized individuals who have a limited ability to communicate thirst. **Communities can potentially prevent hospitalizations by encouraging attention to the fluid status of individuals at risk.**

Urinary Tract Infection (UTI) is usually caused when bacteria enter the bladder and cause inflammation and infection. It is a common condition, with older adults at highest risk. In most cases, an uncomplicated UTI can be treated with proper antibiotics. **Communities can potentially prevent hospitalizations by encouraging individuals to practice good personal hygiene; drink plenty of fluids; and (if practical) avoid conducting urine cultures in asymptomatic patients who have indwelling urethral catheters.**

Angina (without procedures) is chest pain that occurs when a blockage of a coronary artery prevents sufficient oxygen-rich blood from reaching the heart muscle. **Communities can potentially prevent hospitalizations by encouraging regular physical activity; smoking cessation; controlling diabetes, high blood pressure, and abnormal cholesterol; maintaining appropriate body weight; and daily administration of an anti-platelet medication (like low dose aspirin) in most individuals with known coronary artery disease.**

Congestive Heart Failure is the inability of the heart muscle to function well enough to meet the demands of the rest of the body. **Communities can potentially prevent hospitalizations by encouraging individuals to reduce risk factors such as coronary artery disease, diabetes, high cholesterol, high blood pressure, smoking, alcohol abuse, and use of illegal drugs.**

Hypertension (High Blood Pressure) is a syndrome with multiple causes. Hypertension is often controllable with medications. **Communities can potentially prevent hospitalizations by encouraging an increased level of aerobic physical activity, maintaining a healthy weight, limiting the consumption of alcohol to moderate levels for those who drink, reducing salt and sodium intake, and eating a reduced-fat diet high in fruits, vegetables, and low-fat dairy food.**

Asthma occurs when air passages of the lungs become inflamed and narrowed and breathing becomes difficult. Asthma is treatable, and most flare-ups and deaths can be prevented through the use of medications. **Communities can potentially prevent hospitalizations by encouraging people to learn how to recognize particular warning signs of asthma attacks. Treating symptoms early can result in prevented or less severe attacks.**

Chronic Obstructive Pulmonary Disease is characterized by decreased flow in the airways of the lungs. It consists of three related diseases: asthma, chronic bronchitis and emphysema. Because existing medications cannot change the progressive decline in lung function, the goal of medications is to lessen symptoms and/or decrease complications. **Communities can potentially prevent hospitalizations by encouraging education on smoking cessation and minimizing shortness of breath.**

Diabetes Short-term Complications are extreme fluctuations in blood sugar levels. Extreme dizziness and fainting can indicate hypoglycemia (low blood sugar) or hyperglycemia (high blood sugar), and if not brought under control, seizures, shock or coma can occur. Diabetics need to monitor their blood sugar levels carefully and adjust their diet and/or medications accordingly. **Communities can potentially prevent hospitalizations by encouraging the regular monitoring and managing of diabetes in the outpatient health care setting and encouraging patient compliance with treatment plans.**

Diabetes Long-term Complications include risk of developing damage to the eyes, kidneys and nerves. Risk also includes developing cardiovascular disease, including coronary heart disease, stroke, and peripheral vascular disease. Long-term diabetes complications are thought to result from long-term poor control of diabetes. **Communities can potentially prevent hospitalizations by encouraging the regular monitoring and managing of diabetes in the outpatient health care setting and encouraging patient compliance with treatment plans.**

For more information on potentially preventable hospitalizations, go to: www.dshs.state.tx.us/ph.