

Texas Birth Defects Registry (TBDR) Annual Report

Table 2. Prevalence of Selected Birth Defects by Year, Texas, 1999–2020

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|----------------------------------|------|---------------|-------------------|--|
| Central Nervous System | | | | |
| Anencephaly | 1999 | 123 | 3.52 | 2.90 – 4.15 |
| | 2000 | 100 | 2.75 | 2.24 – 3.35 |
| | 2001 | 94 | 2.57 | 2.08 – 3.15 |
| | 2002 | 98 | 2.63 | 2.14 – 3.21 |
| | 2003 | 98 | 2.60 | 2.11 – 3.16 |
| | 2004 | 87 | 2.28 | 1.83 – 2.81 |
| | 2005 | 85 | 2.20 | 1.76 – 2.73 |
| | 2006 | 90 | 2.25 | 1.81 – 2.77 |
| | 2007 | 93 | 2.28 | 1.84 – 2.80 |
| | 2008 | 106 | 2.62 | 2.12 – 3.11 |
| | 2009 | 122 | 3.04 | 2.50 – 3.58 |
| | 2010 | 108 | 2.80 | 2.27 – 3.33 |
| | 2011 | 98 | 2.60 | 2.11 – 3.17 |
| | 2012 | 109 | 2.85 | 2.32 – 3.39 |
| | 2013 | 89 | 2.30 | 1.85 – 2.83 |
| | 2014 | 84 | 2.10 | 1.68 – 2.60 |
| | 2015 | 81 | 2.01 | 1.59 – 2.50 |
| | 2016 | 103 | 2.59 | 2.09 – 3.10 |
| | 2017 | 90 | 2.36 | 1.90 – 2.90 |
| | 2018 | 93 | 2.47 | 1.99 – 3.03 |
| 2019 | 84 | 2.22 | 1.77 – 2.75 | |
| 2020 | 71 | 1.93 | 1.51 – 2.43 | |
| Spina bifida without anencephaly | 1999 | 148 | 4.24 | 3.56 – 4.92 |
| | 2000 | 137 | 3.77 | 3.14 – 4.40 |
| | 2001 | 123 | 3.37 | 2.77 – 3.96 |
| | 2002 | 112 | 3.01 | 2.45 – 3.56 |
| | 2003 | 123 | 3.26 | 2.68 – 3.84 |
| | 2004 | 173 | 4.54 | 3.86 – 5.21 |
| | 2005 | 141 | 3.66 | 3.05 – 4.26 |
| | 2006 | 136 | 3.41 | 2.83 – 3.98 |
| | 2007 | 154 | 3.78 | 3.18 – 4.38 |
| | 2008 | 143 | 3.53 | 2.95 – 4.11 |
| | 2009 | 152 | 3.78 | 3.18 – 4.39 |
| | 2010 | 160 | 4.15 | 3.51 – 4.79 |
| | 2011 | 161 | 4.27 | 3.61 – 4.93 |
| | 2012 | 159 | 4.16 | 3.51 – 4.80 |
| | 2013 | 148 | 3.82 | 3.21 – 4.44 |
| 2014 | 146 | 3.65 | 3.06 – 4.25 | |
| 2015 | 136 | 3.37 | 2.80 – 3.94 | |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|---|------|---------------|-------------------|--|
| | 2016 | 161 | 4.06 | 3.43 – 4.68 |
| | 2017 | 132 | 3.46 | 2.87 – 4.05 |
| | 2018 | 135 | 3.59 | 2.98 – 4.19 |
| | 2019 | 127 | 3.36 | 2.78 – 3.95 |
| | 2020 | 131 | 3.56 | 2.95 – 4.17 |
| Encephalocele | 1999 | 33 | 0.95 | 0.65 – 1.33 |
| | 2000 | 39 | 1.07 | 0.76 – 1.47 |
| | 2001 | 34 | 0.93 | 0.64 – 1.30 |
| | 2002 | 28 | 0.75 | 0.50 – 1.09 |
| | 2003 | 33 | 0.87 | 0.60 – 1.23 |
| | 2004 | 23 | 0.60 | 0.38 – 0.90 |
| | 2005 | 32 | 0.83 | 0.57 – 1.17 |
| | 2006 | 39 | 0.98 | 0.69 – 1.34 |
| | 2007 | 42 | 1.03 | 0.74 – 1.39 |
| | 2008 | 41 | 1.01 | 0.73 – 1.37 |
| | 2009 | 38 | 0.95 | 0.67 – 1.30 |
| | 2010 | 41 | 1.06 | 0.76 – 1.44 |
| | 2011 | 41 | 1.09 | 0.78 – 1.47 |
| | 2012 | 46 | 1.20 | 0.88 – 1.60 |
| | 2013 | 37 | 0.96 | 0.67 – 1.32 |
| | 2014 | 32 | 0.80 | 0.55 – 1.13 |
| | 2015 | 36 | 0.89 | 0.62 – 1.24 |
| | 2016 | 40 | 1.01 | 0.72 – 1.37 |
| | 2017 | 35 | 0.92 | 0.64 – 1.27 |
| | 2018 | 43 | 1.14 | 0.83 – 1.54 |
| | 2019 | 45 | 1.19 | 0.87 – 1.59 |
| | 2020 | 36 | 0.98 | 0.68 – 1.35 |
| Microcephaly, severe (head circumference <3 rd percentile) | 1999 | 94 | 2.69 | 2.18 – 3.29 |
| | 2000 | 118 | 3.25 | 2.66 – 3.83 |
| | 2001 | 92 | 2.52 | 2.03 – 3.09 |
| | 2002 | 119 | 3.20 | 2.62 – 3.77 |
| | 2003 | 104 | 2.76 | 2.23 – 3.29 |
| | 2004 | 131 | 3.43 | 2.85 – 4.02 |
| | 2005 | 153 | 3.97 | 3.34 – 4.60 |
| | 2006 | 152 | 3.81 | 3.20 – 4.41 |
| | 2007 | 140 | 3.44 | 2.87 – 4.01 |
| | 2008 | 177 | 4.37 | 3.72 – 5.01 |
| | 2009 | 180 | 4.48 | 3.83 – 5.14 |
| | 2010 | 197 | 5.11 | 4.39 – 5.82 |
| | 2011 | 178 | 4.72 | 4.02 – 5.41 |
| | 2012 | 218 | 5.70 | 4.94 – 6.46 |
| | 2013 | 234 | 6.04 | 5.27 – 6.82 |
| | 2014 | 217 | 5.43 | 4.71 – 6.15 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|-----------------------------------|------|------------------|----------------------|---|
| | 2015 | 244 | 6.05 | 5.29 – 6.81 |
| | 2016 | 310 | 7.81 | 6.94 – 8.68 |
| | 2017 | 246 | 6.44 | 5.64 – 7.25 |
| | 2018 | 305 | 8.10 | 7.19 – 9.01 |
| | 2019 | 322 | 8.53 | 7.59 – 9.46 |
| | 2020 | 356 | 9.67 | 8.66 – 10.67 |
| Holoprosencephaly | 1999 | 41 | 1.17 | 0.84 – 1.59 |
| | 2000 | 38 | 1.05 | 0.74 – 1.44 |
| | 2001 | 46 | 1.26 | 0.92 – 1.68 |
| | 2002 | 51 | 1.37 | 1.02 – 1.80 |
| | 2003 | 45 | 1.19 | 0.87 – 1.60 |
| | 2004 | 37 | 0.97 | 0.68 – 1.34 |
| | 2005 | 41 | 1.06 | 0.76 – 1.44 |
| | 2006 | 46 | 1.15 | 0.84 – 1.54 |
| | 2007 | 54 | 1.33 | 1.00 – 1.73 |
| | 2008 | 38 | 0.94 | 0.66 – 1.29 |
| | 2009 | 31 | 0.77 | 0.52 – 1.10 |
| | 2010 | 36 | 0.93 | 0.65 – 1.29 |
| | 2011 | 35 | 0.93 | 0.65 – 1.29 |
| | 2012 | 43 | 1.12 | 0.81 – 1.51 |
| | 2013 | 37 | 0.96 | 0.67 – 1.32 |
| | 2014 | 41 | 1.03 | 0.74 – 1.39 |
| | 2015 | 37 | 0.92 | 0.65 – 1.26 |
| | 2016 | 38 | 0.96 | 0.68 – 1.31 |
| | 2017 | 32 | 0.84 | 0.57 – 1.18 |
| | 2018 | 44 | 1.17 | 0.85 – 1.57 |
| | 2019 | 36 | 0.95 | 0.67 – 1.32 |
| | 2020 | 38 | 1.03 | 0.73 – 1.42 |
| Hydrocephaly without spina bifida | 1999 | 295 | 8.45 | 7.48 – 9.41 |
| | 2000 | 264 | 7.27 | 6.39 – 8.14 |
| | 2001 | 234 | 6.41 | 5.59 – 7.23 |
| | 2002 | 251 | 6.74 | 5.91 – 7.57 |
| | 2003 | 249 | 6.60 | 5.78 – 7.42 |
| | 2004 | 265 | 6.95 | 6.11 – 7.78 |
| | 2005 | 234 | 6.07 | 5.29 – 6.85 |
| | 2006 | 270 | 6.76 | 5.96 – 7.57 |
| | 2007 | 309 | 7.58 | 6.74 – 8.43 |
| | 2008 | 296 | 7.30 | 6.47 – 8.14 |
| | 2009 | 302 | 7.52 | 6.67 – 8.37 |
| | 2010 | 298 | 7.73 | 6.85 – 8.60 |
| | 2011 | 311 | 8.24 | 7.33 – 9.16 |
| | 2012 | 339 | 8.86 | 7.92 – 9.81 |
| | 2013 | 330 | 8.52 | 7.60 – 9.44 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|----------------------------|------|---------------|-------------------|--|
| | 2014 | 360 | 9.01 | 8.08 – 9.94 |
| | 2015 | 338 | 8.38 | 7.48 – 9.27 |
| | 2016 | 376 | 9.47 | 8.51 – 10.43 |
| | 2017 | 388 | 10.16 | 9.15 – 11.17 |
| | 2018 | 368 | 9.77 | 8.78 – 10.77 |
| | 2019 | 357 | 9.45 | 8.47 – 10.43 |
| | 2020 | 368 | 9.99 | 8.97 – 11.01 |
| Eye and Ear | | | | |
| Anophthalmia | 1999 | 12 | 0.34 | 0.18 – 0.60 |
| | 2000 | 11 | 0.30 | 0.15 – 0.54 |
| | 2001 | 12 | 0.33 | 0.17 – 0.57 |
| | 2002 | 11 | 0.30 | 0.15 – 0.53 |
| | 2003 | 12 | 0.32 | 0.16 – 0.56 |
| | 2004 | 10 | 0.26 | 0.13 – 0.48 |
| | 2005 | 14 | 0.36 | 0.20 – 0.61 |
| | 2006 | 17 | 0.43 | 0.25 – 0.68 |
| | 2007 | 13 | 0.32 | 0.17 – 0.55 |
| | 2008 | 9 | 0.22 | 0.10 – 0.42 |
| | 2009 | 13 | 0.32 | 0.17 – 0.55 |
| | 2010 | 13 | 0.34 | 0.18 – 0.58 |
| | 2011 | 19 | 0.50 | 0.30 – 0.79 |
| | 2012 | 7 | 0.18 | 0.07 – 0.38 |
| | 2013 | 10 | 0.26 | 0.12 – 0.48 |
| | 2014 | 6 | 0.15 | 0.06 – 0.33 |
| | 2015 | 8 | 0.20 | 0.09 – 0.39 |
| | 2016 | 9 | 0.23 | 0.10 – 0.43 |
| | 2017 | 8 | 0.21 | 0.09 – 0.41 |
| | 2018 | 11 | 0.29 | 0.15 – 0.52 |
| | 2019 | 9 | 0.24 | 0.11 – 0.45 |
| | 2020 | 12 | 0.33 | 0.17 – 0.57 |
| Microphthalmia | 1999 | 85 | 2.43 | 1.94 – 3.01 |
| | 2000 | 91 | 2.50 | 2.02 – 3.08 |
| | 2001 | 89 | 2.44 | 1.96 – 3.00 |
| | 2002 | 106 | 2.85 | 2.30 – 3.39 |
| | 2003 | 102 | 2.70 | 2.18 – 3.23 |
| | 2004 | 100 | 2.62 | 2.13 – 3.19 |
| | 2005 | 105 | 2.72 | 2.20 – 3.24 |
| | 2006 | 120 | 3.01 | 2.47 – 3.54 |
| | 2007 | 131 | 3.22 | 2.66 – 3.77 |
| | 2008 | 120 | 2.96 | 2.43 – 3.49 |
| | 2009 | 104 | 2.59 | 2.09 – 3.09 |
| | 2010 | 115 | 2.98 | 2.44 – 3.53 |
| | 2011 | 119 | 3.15 | 2.59 – 3.72 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|----------------------------|------|------------------|----------------------|---|
| | 2012 | 100 | 2.61 | 2.13 – 3.18 |
| | 2013 | 101 | 2.61 | 2.10 – 3.12 |
| | 2014 | 101 | 2.53 | 2.04 – 3.02 |
| | 2015 | 94 | 2.33 | 1.88 – 2.85 |
| | 2016 | 117 | 2.95 | 2.41 – 3.48 |
| | 2017 | 101 | 2.64 | 2.13 – 3.16 |
| | 2018 | 103 | 2.74 | 2.21 – 3.26 |
| | 2019 | 109 | 2.89 | 2.34 – 3.43 |
| | 2020 | 95 | 2.58 | 2.09 – 3.15 |
| Cataract | 1999 | 42 | 1.20 | 0.87 – 1.63 |
| | 2000 | 63 | 1.73 | 1.33 – 2.22 |
| | 2001 | 54 | 1.48 | 1.11 – 1.93 |
| | 2002 | 75 | 2.01 | 1.58 – 2.52 |
| | 2003 | 65 | 1.72 | 1.33 – 2.20 |
| | 2004 | 72 | 1.89 | 1.48 – 2.38 |
| | 2005 | 78 | 2.02 | 1.60 – 2.52 |
| | 2006 | 81 | 2.03 | 1.61 – 2.52 |
| | 2007 | 93 | 2.28 | 1.84 – 2.80 |
| | 2008 | 73 | 1.80 | 1.41 – 2.26 |
| | 2009 | 75 | 1.87 | 1.47 – 2.34 |
| | 2010 | 80 | 2.07 | 1.64 – 2.58 |
| | 2011 | 85 | 2.25 | 1.80 – 2.79 |
| | 2012 | 63 | 1.65 | 1.27 – 2.11 |
| | 2013 | 64 | 1.65 | 1.27 – 2.11 |
| | 2014 | 76 | 1.90 | 1.50 – 2.38 |
| | 2015 | 78 | 1.93 | 1.53 – 2.41 |
| | 2016 | 99 | 2.49 | 2.03 – 3.04 |
| | 2017 | 80 | 2.09 | 1.66 – 2.61 |
| | 2018 | 83 | 2.20 | 1.76 – 2.73 |
| | 2019 | 80 | 2.12 | 1.68 – 2.64 |
| | 2020 | 71 | 1.93 | 1.51 – 2.43 |
| Anotia or microtia | 1999 | 100 | 2.86 | 2.33 – 3.48 |
| | 2000 | 101 | 2.78 | 2.24 – 3.32 |
| | 2001 | 106 | 2.90 | 2.35 – 3.46 |
| | 2002 | 107 | 2.87 | 2.33 – 3.42 |
| | 2003 | 102 | 2.70 | 2.18 – 3.23 |
| | 2004 | 130 | 3.41 | 2.82 – 3.99 |
| | 2005 | 116 | 3.01 | 2.46 – 3.56 |
| | 2006 | 134 | 3.36 | 2.79 – 3.92 |
| | 2007 | 126 | 3.09 | 2.55 – 3.63 |
| | 2008 | 164 | 4.05 | 3.43 – 4.67 |
| | 2009 | 138 | 3.44 | 2.86 – 4.01 |
| | 2010 | 143 | 3.71 | 3.10 – 4.31 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|------------------------------------|------|---------------|-------------------|--|
| | 2011 | 136 | 3.60 | 3.00 – 4.21 |
| | 2012 | 128 | 3.35 | 2.77 – 3.93 |
| | 2013 | 139 | 3.59 | 2.99 – 4.19 |
| | 2014 | 147 | 3.68 | 3.08 – 4.27 |
| | 2015 | 161 | 3.99 | 3.37 – 4.61 |
| | 2016 | 171 | 4.31 | 3.66 – 4.95 |
| | 2017 | 155 | 4.06 | 3.42 – 4.70 |
| | 2018 | 161 | 4.28 | 3.62 – 4.94 |
| | 2019 | 166 | 4.39 | 3.73 – 5.06 |
| | 2020 | 172 | 4.67 | 3.97 – 5.37 |
| Cardiac and Circulatory | | | | |
| Common truncus | 1999 | 17 | 0.49 | 0.28 – 0.78 |
| | 2000 | 27 | 0.74 | 0.49 – 1.08 |
| | 2001 | 26 | 0.71 | 0.47 – 1.04 |
| | 2002 | 29 | 0.78 | 0.52 – 1.12 |
| | 2003 | 28 | 0.74 | 0.49 – 1.07 |
| | 2004 | 31 | 0.81 | 0.55 – 1.15 |
| | 2005 | 24 | 0.62 | 0.40 – 0.93 |
| | 2006 | 28 | 0.70 | 0.47 – 1.01 |
| | 2007 | 29 | 0.71 | 0.48 – 1.02 |
| | 2008 | 44 | 1.09 | 0.79 – 1.46 |
| | 2009 | 24 | 0.60 | 0.38 – 0.89 |
| | 2010 | 31 | 0.80 | 0.55 – 1.14 |
| | 2011 | 30 | 0.80 | 0.54 – 1.14 |
| | 2012 | 39 | 1.02 | 0.73 – 1.39 |
| | 2013 | 34 | 0.88 | 0.61 – 1.23 |
| | 2014 | 31 | 0.78 | 0.53 – 1.10 |
| | 2015 | 30 | 0.74 | 0.50 – 1.06 |
| | 2016 | 30 | 0.76 | 0.51 – 1.08 |
| | 2017 | 24 | 0.63 | 0.40 – 0.94 |
| | 2018 | 34 | 0.90 | 0.63 – 1.26 |
| | 2019 | 28 | 0.74 | 0.49 – 1.07 |
| | 2020 | 27 | 0.73 | 0.48 – 1.07 |
| Transposition of the great vessels | 1999 | 132 | 3.78 | 3.14 – 4.43 |
| | 2000 | 118 | 3.25 | 2.66 – 3.83 |
| | 2001 | 138 | 3.78 | 3.15 – 4.41 |
| | 2002 | 103 | 2.77 | 2.23 – 3.30 |
| | 2003 | 131 | 3.47 | 2.88 – 4.07 |
| | 2004 | 128 | 3.36 | 2.77 – 3.94 |
| | 2005 | 123 | 3.19 | 2.63 – 3.75 |
| | 2006 | 149 | 3.73 | 3.13 – 4.33 |
| | 2007 | 147 | 3.61 | 3.02 – 4.19 |
| | 2008 | 151 | 3.73 | 3.13 – 4.32 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|-------------------------------|------|---------------|-------------------|--|
| | 2009 | 148 | 3.69 | 3.09 – 4.28 |
| | 2010 | 150 | 3.89 | 3.27 – 4.51 |
| | 2011 | 125 | 3.31 | 2.73 – 3.89 |
| | 2012 | 124 | 3.24 | 2.67 – 3.81 |
| | 2013 | 150 | 3.87 | 3.25 – 4.49 |
| | 2014 | 142 | 3.55 | 2.97 – 4.14 |
| | 2015 | 127 | 3.15 | 2.60 – 3.70 |
| | 2016 | 117 | 2.95 | 2.41 – 3.48 |
| | 2017 | 103 | 2.70 | 2.18 – 3.22 |
| | 2018 | 119 | 3.16 | 2.59 – 3.73 |
| | 2019 | 105 | 2.78 | 2.25 – 3.31 |
| | 2020 | 112 | 3.04 | 2.48 – 3.60 |
| Double outlet right ventricle | 1999 | 65 | 1.86 | 1.44 – 2.37 |
| | 2000 | 58 | 1.60 | 1.21 – 2.06 |
| | 2001 | 82 | 2.25 | 1.79 – 2.79 |
| | 2002 | 74 | 1.99 | 1.56 – 2.49 |
| | 2003 | 78 | 2.07 | 1.63 – 2.58 |
| | 2004 | 76 | 1.99 | 1.57 – 2.49 |
| | 2005 | 76 | 1.97 | 1.55 – 2.47 |
| | 2006 | 74 | 1.85 | 1.46 – 2.33 |
| | 2007 | 86 | 2.11 | 1.69 – 2.61 |
| | 2008 | 102 | 2.52 | 2.03 – 3.01 |
| | 2009 | 107 | 2.66 | 2.16 – 3.17 |
| | 2010 | 81 | 2.10 | 1.67 – 2.61 |
| | 2011 | 91 | 2.41 | 1.94 – 2.96 |
| | 2012 | 85 | 2.22 | 1.78 – 2.75 |
| | 2013 | 120 | 3.10 | 2.55 – 3.65 |
| | 2014 | 98 | 2.45 | 1.99 – 2.99 |
| | 2015 | 90 | 2.23 | 1.79 – 2.74 |
| | 2016 | 103 | 2.59 | 2.09 – 3.10 |
| | 2017 | 86 | 2.25 | 1.80 – 2.78 |
| | 2018 | 97 | 2.58 | 2.09 – 3.14 |
| | 2019 | 112 | 2.97 | 2.42 – 3.51 |
| | 2020 | 107 | 2.91 | 2.35 – 3.46 |
| Tetralogy of Fallot | 1999 | 110 | 3.15 | 2.56 – 3.74 |
| | 2000 | 104 | 2.86 | 2.31 – 3.41 |
| | 2001 | 127 | 3.48 | 2.87 – 4.08 |
| | 2002 | 132 | 3.54 | 2.94 – 4.15 |
| | 2003 | 144 | 3.82 | 3.19 – 4.44 |
| | 2004 | 110 | 2.88 | 2.34 – 3.42 |
| | 2005 | 152 | 3.94 | 3.32 – 4.57 |
| | 2006 | 152 | 3.81 | 3.20 – 4.41 |
| | 2007 | 136 | 3.34 | 2.78 – 3.90 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|----------------------------|------|------------------|----------------------|---|
| | 2008 | 180 | 4.44 | 3.79 – 5.09 |
| | 2009 | 149 | 3.71 | 3.11 – 4.31 |
| | 2010 | 141 | 3.66 | 3.05 – 4.26 |
| | 2011 | 164 | 4.35 | 3.68 – 5.01 |
| | 2012 | 179 | 4.68 | 3.99 – 5.37 |
| | 2013 | 155 | 4.00 | 3.37 – 4.63 |
| | 2014 | 183 | 4.58 | 3.92 – 5.24 |
| | 2015 | 173 | 4.29 | 3.65 – 4.93 |
| | 2016 | 198 | 4.99 | 4.29 – 5.68 |
| | 2017 | 161 | 4.22 | 3.56 – 4.87 |
| | 2018 | 173 | 4.59 | 3.91 – 5.28 |
| | 2019 | 170 | 4.50 | 3.82 – 5.18 |
| | 2020 | 144 | 3.91 | 3.27 – 4.55 |
| Ventricular septal defect | 1999 | 1,425 | 40.81 | 38.69 – 42.93 |
| | 2000 | 1,557 | 42.85 | 40.73 – 44.98 |
| | 2001 | 1,673 | 45.82 | 43.63 – 48.02 |
| | 2002 | 1,823 | 48.96 | 46.71 – 51.20 |
| | 2003 | 1,847 | 48.94 | 46.71 – 51.18 |
| | 2004 | 1,969 | 51.62 | 49.34 – 53.90 |
| | 2005 | 2,205 | 57.19 | 54.81 – 59.58 |
| | 2006 | 2,275 | 56.97 | 54.63 – 59.31 |
| | 2007 | 2,485 | 60.99 | 58.59 – 63.39 |
| | 2008 | 2,624 | 64.75 | 62.27 – 67.23 |
| | 2009 | 2,707 | 67.41 | 64.87 – 69.94 |
| | 2010 | 2,607 | 67.58 | 64.99 – 70.18 |
| | 2011 | 2,551 | 67.62 | 64.99 – 70.24 |
| | 2012 | 2,520 | 65.89 | 63.32 – 68.47 |
| | 2013 | 2,514 | 64.94 | 62.40 – 67.48 |
| | 2014 | 2,629 | 65.81 | 63.29 – 68.33 |
| | 2015 | 2,785 | 69.03 | 66.47 – 71.60 |
| | 2016 | 2,711 | 68.29 | 65.72 – 70.86 |
| | 2017 | 2,672 | 69.97 | 67.32 – 72.62 |
| | 2018 | 2,681 | 71.21 | 68.51 – 73.90 |
| | 2019 | 2,642 | 69.95 | 67.28 – 72.62 |
| | 2020 | 2,645 | 71.81 | 69.08 – 74.55 |
| Atrial septal defect | 1999 | 1,377 | 39.44 | 37.35 – 41.52 |
| | 2000 | 1,424 | 39.19 | 37.16 – 41.23 |
| | 2001 | 1,523 | 41.72 | 39.62 – 43.81 |
| | 2002 | 1,621 | 43.53 | 41.41 – 45.65 |
| | 2003 | 1,809 | 47.94 | 45.73 – 50.15 |
| | 2004 | 2,135 | 55.97 | 53.60 – 58.35 |
| | 2005 | 2,500 | 64.84 | 62.30 – 67.39 |
| | 2006 | 2,469 | 61.83 | 59.39 – 64.27 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|---|------|---------------|-------------------|--|
| | 2007 | 2,463 | 60.45 | 58.06 – 62.84 |
| | 2008 | 2,808 | 69.29 | 66.73 – 71.85 |
| | 2009 | 2,707 | 67.41 | 64.87 – 69.94 |
| | 2010 | 2,962 | 76.79 | 74.02 – 79.55 |
| | 2011 | 3,259 | 86.38 | 83.42 – 89.35 |
| | 2012 | 3,223 | 84.28 | 81.37 – 87.18 |
| | 2013 | 3,247 | 83.88 | 80.99 – 86.76 |
| | 2014 | 3,432 | 85.91 | 83.04 – 88.79 |
| | 2015 | 3,655 | 90.60 | 87.66 – 93.53 |
| | 2016 | 3,716 | 93.60 | 90.59 – 96.61 |
| | 2017 | 3,350 | 87.72 | 84.75 – 90.70 |
| | 2018 | 3,326 | 88.34 | 85.34 – 91.34 |
| | 2019 | 3,404 | 90.12 | 87.09 – 93.15 |
| | 2020 | 3,586 | 97.36 | 94.18 – 100.55 |
| Atrioventricular septal defect (endocardial cushion defect) | 1999 | 138 | 3.95 | 3.29 – 4.61 |
| | 2000 | 152 | 4.18 | 3.52 – 4.85 |
| | 2001 | 159 | 4.36 | 3.68 – 5.03 |
| | 2002 | 151 | 4.06 | 3.41 – 4.70 |
| | 2003 | 145 | 3.84 | 3.22 – 4.47 |
| | 2004 | 152 | 3.98 | 3.35 – 4.62 |
| | 2005 | 155 | 4.02 | 3.39 – 4.65 |
| | 2006 | 181 | 4.53 | 3.87 – 5.19 |
| | 2007 | 157 | 3.85 | 3.25 – 4.46 |
| | 2008 | 198 | 4.89 | 4.21 – 5.57 |
| | 2009 | 184 | 4.58 | 3.92 – 5.24 |
| | 2010 | 167 | 4.33 | 3.67 – 4.99 |
| | 2011 | 163 | 4.32 | 3.66 – 4.98 |
| | 2012 | 166 | 4.34 | 3.68 – 5.00 |
| | 2013 | 191 | 4.93 | 4.23 – 5.63 |
| | 2014 | 187 | 4.68 | 4.01 – 5.35 |
| | 2015 | 184 | 4.56 | 3.90 – 5.22 |
| | 2016 | 182 | 4.58 | 3.92 – 5.25 |
| | 2017 | 186 | 4.87 | 4.17 – 5.57 |
| | 2018 | 168 | 4.46 | 3.79 – 5.14 |
| | 2019 | 152 | 4.02 | 3.38 – 4.66 |
| | 2020 | 163 | 4.43 | 3.75 – 5.10 |
| Pulmonary valve atresia or stenosis | 1999 | 220 | 6.30 | 5.47 – 7.13 |
| | 2000 | 235 | 6.47 | 5.64 – 7.30 |
| | 2001 | 272 | 7.45 | 6.56 – 8.34 |
| | 2002 | 249 | 6.69 | 5.86 – 7.52 |
| | 2003 | 319 | 8.45 | 7.53 – 9.38 |
| | 2004 | 344 | 9.02 | 8.07 – 9.97 |
| | 2005 | 346 | 8.97 | 8.03 – 9.92 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|-------------------------------------|------|---------------|-------------------|--|
| | 2006 | 347 | 8.69 | 7.78 – 9.60 |
| | 2007 | 403 | 9.89 | 8.93 – 10.86 |
| | 2008 | 467 | 11.52 | 10.48 – 12.57 |
| | 2009 | 416 | 10.36 | 9.36 – 11.35 |
| | 2010 | 429 | 11.12 | 10.07 – 12.17 |
| | 2011 | 464 | 12.30 | 11.18 – 13.42 |
| | 2012 | 437 | 11.43 | 10.36 – 12.50 |
| | 2013 | 410 | 10.59 | 9.57 – 11.62 |
| | 2014 | 433 | 10.84 | 9.82 – 11.86 |
| | 2015 | 403 | 9.99 | 9.01 – 10.96 |
| | 2016 | 425 | 10.71 | 9.69 – 11.72 |
| | 2017 | 449 | 11.76 | 10.67 – 12.85 |
| | 2018 | 455 | 12.08 | 10.97 – 13.20 |
| | 2019 | 422 | 11.17 | 10.11 – 12.24 |
| | 2020 | 455 | 12.35 | 11.22 – 13.49 |
| Tricuspid valve atresia or stenosis | 1999 | 58 | 1.66 | 1.26 – 2.15 |
| | 2000 | 41 | 1.13 | 0.81 – 1.53 |
| | 2001 | 57 | 1.56 | 1.18 – 2.02 |
| | 2002 | 55 | 1.48 | 1.11 – 1.92 |
| | 2003 | 65 | 1.72 | 1.33 – 2.20 |
| | 2004 | 68 | 1.78 | 1.38 – 2.26 |
| | 2005 | 60 | 1.56 | 1.19 – 2.00 |
| | 2006 | 73 | 1.83 | 1.43 – 2.30 |
| | 2007 | 73 | 1.79 | 1.40 – 2.25 |
| | 2008 | 68 | 1.68 | 1.30 – 2.13 |
| | 2009 | 64 | 1.59 | 1.23 – 2.04 |
| | 2010 | 79 | 2.05 | 1.62 – 2.55 |
| | 2011 | 75 | 1.99 | 1.56 – 2.49 |
| | 2012 | 64 | 1.67 | 1.29 – 2.14 |
| | 2013 | 78 | 2.01 | 1.59 – 2.51 |
| | 2014 | 88 | 2.20 | 1.77 – 2.71 |
| | 2015 | 84 | 2.08 | 1.66 – 2.58 |
| | 2016 | 72 | 1.81 | 1.42 – 2.28 |
| | 2017 | 106 | 2.78 | 2.25 – 3.30 |
| | 2018 | 95 | 2.52 | 2.04 – 3.08 |
| | 2019 | 73 | 1.93 | 1.51 – 2.43 |
| | 2020 | 95 | 2.58 | 2.09 – 3.15 |
| Ebstein anomaly | 1999 | 24 | 0.69 | 0.44 – 1.02 |
| | 2000 | 27 | 0.74 | 0.49 – 1.08 |
| | 2001 | 19 | 0.52 | 0.31 – 0.81 |
| | 2002 | 23 | 0.62 | 0.39 – 0.93 |
| | 2003 | 31 | 0.82 | 0.56 – 1.17 |
| | 2004 | 41 | 1.07 | 0.77 – 1.46 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|---------------------------------|------|---------------|-------------------|--|
| | 2005 | 23 | 0.60 | 0.38 – 0.90 |
| | 2006 | 25 | 0.63 | 0.41 – 0.92 |
| | 2007 | 24 | 0.59 | 0.38 – 0.88 |
| | 2008 | 29 | 0.72 | 0.48 – 1.03 |
| | 2009 | 23 | 0.57 | 0.36 – 0.86 |
| | 2010 | 30 | 0.78 | 0.52 – 1.11 |
| | 2011 | 31 | 0.82 | 0.56 – 1.17 |
| | 2012 | 29 | 0.76 | 0.51 – 1.09 |
| | 2013 | 33 | 0.85 | 0.59 – 1.20 |
| | 2014 | 34 | 0.85 | 0.59 – 1.19 |
| | 2015 | 28 | 0.69 | 0.46 – 1.00 |
| | 2016 | 22 | 0.55 | 0.35 – 0.84 |
| | 2017 | 39 | 1.02 | 0.73 – 1.40 |
| | 2018 | 39 | 1.04 | 0.74 – 1.42 |
| | 2019 | 39 | 1.03 | 0.73 – 1.41 |
| | 2020 | 26 | 0.71 | 0.46 – 1.03 |
| Aortic valve stenosis | 1999 | 85 | 2.43 | 1.94 – 3.01 |
| | 2000 | 77 | 2.12 | 1.67 – 2.65 |
| | 2001 | 84 | 2.30 | 1.84 – 2.85 |
| | 2002 | 90 | 2.42 | 1.94 – 2.97 |
| | 2003 | 95 | 2.52 | 2.04 – 3.08 |
| | 2004 | 99 | 2.60 | 2.11 – 3.16 |
| | 2005 | 76 | 1.97 | 1.55 – 2.47 |
| | 2006 | 92 | 2.30 | 1.86 – 2.83 |
| | 2007 | 89 | 2.18 | 1.75 – 2.69 |
| | 2008 | 110 | 2.71 | 2.21 – 3.22 |
| | 2009 | 97 | 2.42 | 1.96 – 2.95 |
| | 2010 | 104 | 2.70 | 2.18 – 3.21 |
| | 2011 | 110 | 2.92 | 2.37 – 3.46 |
| | 2012 | 111 | 2.90 | 2.36 – 3.44 |
| | 2013 | 85 | 2.20 | 1.75 – 2.72 |
| | 2014 | 101 | 2.53 | 2.04 – 3.02 |
| | 2015 | 97 | 2.40 | 1.95 – 2.93 |
| | 2016 | 83 | 2.09 | 1.67 – 2.59 |
| | 2017 | 70 | 1.83 | 1.43 – 2.32 |
| | 2018 | 103 | 2.74 | 2.21 – 3.26 |
| | 2019 | 92 | 2.44 | 1.96 – 2.99 |
| | 2020 | 80 | 2.17 | 1.72 – 2.70 |
| Hypoplastic left heart syndrome | 1999 | 66 | 1.89 | 1.46 – 2.40 |
| | 2000 | 81 | 2.23 | 1.77 – 2.77 |
| | 2001 | 67 | 1.84 | 1.42 – 2.33 |
| | 2002 | 90 | 2.42 | 1.94 – 2.97 |
| | 2003 | 74 | 1.96 | 1.54 – 2.46 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|----------------------------|------|------------------|----------------------|---|
| | 2004 | 83 | 2.18 | 1.73 – 2.70 |
| | 2005 | 77 | 2.00 | 1.58 – 2.50 |
| | 2006 | 87 | 2.18 | 1.75 – 2.69 |
| | 2007 | 80 | 1.96 | 1.56 – 2.44 |
| | 2008 | 89 | 2.20 | 1.76 – 2.70 |
| | 2009 | 86 | 2.14 | 1.71 – 2.64 |
| | 2010 | 93 | 2.41 | 1.95 – 2.95 |
| | 2011 | 86 | 2.28 | 1.82 – 2.82 |
| | 2012 | 104 | 2.72 | 2.20 – 3.24 |
| | 2013 | 99 | 2.56 | 2.08 – 3.11 |
| | 2014 | 89 | 2.23 | 1.79 – 2.74 |
| | 2015 | 107 | 2.65 | 2.15 – 3.15 |
| | 2016 | 93 | 2.34 | 1.89 – 2.87 |
| | 2017 | 84 | 2.20 | 1.75 – 2.72 |
| | 2018 | 91 | 2.42 | 1.95 – 2.97 |
| | 2019 | 96 | 2.54 | 2.06 – 3.10 |
| | 2020 | 75 | 2.04 | 1.60 – 2.55 |
| Patent ductus arteriosus | 1999 | 1,509 | 43.22 | 41.04 – 45.40 |
| | 2000 | 1,498 | 41.23 | 39.14 – 43.32 |
| | 2001 | 1,605 | 43.96 | 41.81 – 46.11 |
| | 2002 | 1,641 | 44.07 | 41.94 – 46.20 |
| | 2003 | 1,799 | 47.67 | 45.47 – 49.87 |
| | 2004 | 2,048 | 53.69 | 51.37 – 56.02 |
| | 2005 | 2,011 | 52.16 | 49.88 – 54.44 |
| | 2006 | 2,182 | 54.64 | 52.35 – 56.94 |
| | 2007 | 2,141 | 52.55 | 50.32 – 54.77 |
| | 2008 | 2,513 | 62.01 | 59.59 – 64.44 |
| | 2009 | 2,663 | 66.31 | 63.79 – 68.83 |
| | 2010 | 2,688 | 69.68 | 67.05 – 72.32 |
| | 2011 | 2,686 | 71.19 | 68.50 – 73.89 |
| | 2012 | 2,706 | 70.76 | 68.09 – 73.42 |
| | 2013 | 2,862 | 73.93 | 71.22 – 76.64 |
| | 2014 | 2,886 | 72.24 | 69.61 – 74.88 |
| | 2015 | 2,947 | 73.05 | 70.41 – 75.68 |
| | 2016 | 2,854 | 71.89 | 69.25 – 74.53 |
| | 2017 | 2,516 | 65.89 | 63.31 – 68.46 |
| | 2018 | 2,377 | 63.13 | 60.60 – 65.67 |
| | 2019 | 2,023 | 53.56 | 51.23 – 55.89 |
| | 2020 | 1,118 | 30.35 | 28.57 – 32.13 |
| Coarctation of the aorta | 1999 | 152 | 4.35 | 3.66 – 5.05 |
| | 2000 | 147 | 4.05 | 3.39 – 4.70 |
| | 2001 | 154 | 4.22 | 3.55 – 4.88 |
| | 2002 | 195 | 5.24 | 4.50 – 5.97 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|----------------------------|------|---------------|-------------------|--|
| | 2003 | 196 | 5.19 | 4.47 – 5.92 |
| | 2004 | 179 | 4.69 | 4.01 – 5.38 |
| | 2005 | 201 | 5.21 | 4.49 – 5.93 |
| | 2006 | 223 | 5.58 | 4.85 – 6.32 |
| | 2007 | 193 | 4.74 | 4.07 – 5.41 |
| | 2008 | 210 | 5.18 | 4.48 – 5.88 |
| | 2009 | 205 | 5.10 | 4.41 – 5.80 |
| | 2010 | 211 | 5.47 | 4.73 – 6.21 |
| | 2011 | 210 | 5.57 | 4.81 – 6.32 |
| | 2012 | 219 | 5.73 | 4.97 – 6.48 |
| | 2013 | 233 | 6.02 | 5.25 – 6.79 |
| | 2014 | 239 | 5.98 | 5.22 – 6.74 |
| | 2015 | 193 | 4.78 | 4.11 – 5.46 |
| | 2016 | 215 | 5.42 | 4.69 – 6.14 |
| | 2017 | 226 | 5.92 | 5.15 – 6.69 |
| | 2018 | 251 | 6.67 | 5.84 – 7.49 |
| | 2019 | 213 | 5.64 | 4.88 – 6.40 |
| | 2020 | 203 | 5.51 | 4.75 – 6.27 |

Respiratory

| | | | | |
|--|------|------|-------------|-------------|
| Choanal atresia or stenosis | 1999 | 43 | 1.23 | 0.89 – 1.66 |
| | 2000 | 48 | 1.32 | 0.97 – 1.75 |
| | 2001 | 42 | 1.15 | 0.83 – 1.55 |
| | 2002 | 45 | 1.21 | 0.88 – 1.62 |
| | 2003 | 38 | 1.01 | 0.71 – 1.38 |
| | 2004 | 32 | 0.84 | 0.57 – 1.18 |
| | 2005 | 43 | 1.12 | 0.81 – 1.50 |
| | 2006 | 59 | 1.48 | 1.12 – 1.91 |
| | 2007 | 44 | 1.08 | 0.78 – 1.45 |
| | 2008 | 55 | 1.36 | 1.02 – 1.77 |
| | 2009 | 56 | 1.39 | 1.05 – 1.81 |
| | 2010 | 50 | 1.30 | 0.96 – 1.71 |
| | 2011 | 52 | 1.38 | 1.03 – 1.81 |
| | 2012 | 51 | 1.33 | 0.99 – 1.75 |
| | 2013 | 55 | 1.42 | 1.07 – 1.85 |
| | 2014 | 63 | 1.58 | 1.21 – 2.02 |
| | 2015 | 58 | 1.44 | 1.09 – 1.86 |
| | 2016 | 33 | 0.83 | 0.57 – 1.17 |
| | 2017 | 60 | 1.57 | 1.20 – 2.02 |
| | 2018 | 43 | 1.14 | 0.83 – 1.54 |
| 2019 | 57 | 1.51 | 1.14 – 1.96 | |
| 2020 | 49 | 1.33 | 0.98 – 1.76 | |
| Agenesis, aplasia, or hypoplasia of the lung | 1999 | 186 | 5.33 | 4.56 – 6.09 |
| | 2000 | 157 | 4.32 | 3.65 – 5.00 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|----------------------------|------|---------------|-------------------|--|
| | 2001 | 157 | 4.30 | 3.63 – 4.97 |
| | 2002 | 135 | 3.63 | 3.01 – 4.24 |
| | 2003 | 122 | 3.23 | 2.66 – 3.81 |
| | 2004 | 99 | 2.60 | 2.11 – 3.16 |
| | 2005 | 110 | 2.85 | 2.32 – 3.39 |
| | 2006 | 110 | 2.75 | 2.24 – 3.27 |
| | 2007 | 104 | 2.55 | 2.06 – 3.04 |
| | 2008 | 94 | 2.32 | 1.87 – 2.84 |
| | 2009 | 115 | 2.86 | 2.34 – 3.39 |
| | 2010 | 114 | 2.96 | 2.41 – 3.50 |
| | 2011 | 105 | 2.78 | 2.25 – 3.32 |
| | 2012 | 90 | 2.35 | 1.89 – 2.89 |
| | 2013 | 117 | 3.02 | 2.47 – 3.57 |
| | 2014 | 103 | 2.58 | 2.08 – 3.08 |
| | 2015 | 116 | 2.88 | 2.35 – 3.40 |
| | 2016 | 114 | 2.87 | 2.34 – 3.40 |
| | 2017 | 110 | 2.88 | 2.34 – 3.42 |
| | 2018 | 103 | 2.74 | 2.21 – 3.26 |
| | 2019 | 76 | 2.01 | 1.59 – 2.52 |
| | 2020 | 79 | 2.14 | 1.70 – 2.67 |

Oral Clefts

| | | | | |
|--|------|-----|------|-------------|
| Cleft palate alone (without cleft lip) | 1999 | 213 | 6.10 | 5.28 – 6.92 |
| | 2000 | 227 | 6.25 | 5.44 – 7.06 |
| | 2001 | 216 | 5.92 | 5.13 – 6.71 |
| | 2002 | 196 | 5.26 | 4.53 – 6.00 |
| | 2003 | 202 | 5.35 | 4.61 – 6.09 |
| | 2004 | 210 | 5.51 | 4.76 – 6.25 |
| | 2005 | 254 | 6.59 | 5.78 – 7.40 |
| | 2006 | 222 | 5.56 | 4.83 – 6.29 |
| | 2007 | 239 | 5.87 | 5.12 – 6.61 |
| | 2008 | 264 | 6.51 | 5.73 – 7.30 |
| | 2009 | 257 | 6.40 | 5.62 – 7.18 |
| | 2010 | 271 | 7.03 | 6.19 – 7.86 |
| | 2011 | 211 | 5.59 | 4.84 – 6.35 |
| | 2012 | 197 | 5.15 | 4.43 – 5.87 |
| | 2013 | 238 | 6.15 | 5.37 – 6.93 |
| | 2014 | 221 | 5.53 | 4.80 – 6.26 |
| | 2015 | 243 | 6.02 | 5.27 – 6.78 |
| | 2016 | 227 | 5.72 | 4.97 – 6.46 |
| | 2017 | 226 | 5.92 | 5.15 – 6.69 |
| | 2018 | 251 | 6.67 | 5.84 – 7.49 |
| | 2019 | 213 | 5.64 | 4.88 – 6.40 |
| | 2020 | 215 | 5.84 | 5.06 – 6.62 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|--|------|------------------|----------------------|---|
| Cleft lip with or without cleft palate | 1999 | 372 | 10.65 | 9.57 – 11.74 |
| | 2000 | 401 | 11.04 | 9.96 – 12.12 |
| | 2001 | 401 | 10.98 | 9.91 – 12.06 |
| | 2002 | 389 | 10.45 | 9.41 – 11.48 |
| | 2003 | 390 | 10.33 | 9.31 – 11.36 |
| | 2004 | 452 | 11.85 | 10.76 – 12.94 |
| | 2005 | 430 | 11.15 | 10.10 – 12.21 |
| | 2006 | 447 | 11.19 | 10.16 – 12.23 |
| | 2007 | 434 | 10.65 | 9.65 – 11.65 |
| | 2008 | 441 | 10.88 | 9.87 – 11.90 |
| | 2009 | 378 | 9.41 | 8.46 – 10.36 |
| | 2010 | 394 | 10.21 | 9.21 – 11.22 |
| | 2011 | 454 | 12.03 | 10.93 – 13.14 |
| | 2012 | 409 | 10.69 | 9.66 – 11.73 |
| | 2013 | 388 | 10.02 | 9.03 – 11.02 |
| | 2014 | 437 | 10.94 | 9.91 – 11.96 |
| | 2015 | 445 | 11.03 | 10.01 – 12.06 |
| | 2016 | 413 | 10.40 | 9.40 – 11.41 |
| | 2017 | 375 | 9.82 | 8.83 – 10.81 |
| | 2018 | 455 | 12.08 | 10.97 – 13.20 |
| 2019 | 388 | 10.27 | 9.25 – 11.29 | |
| 2020 | 382 | 10.37 | 9.33 – 11.41 | |
| Gastrointestinal | | | | |
| Tracheoesophageal fistula/ esophageal atresia | 1999 | 78 | 2.23 | 1.77 – 2.79 |
| | 2000 | 72 | 1.98 | 1.55 – 2.50 |
| | 2001 | 72 | 1.97 | 1.54 – 2.48 |
| | 2002 | 86 | 2.31 | 1.85 – 2.85 |
| | 2003 | 78 | 2.07 | 1.63 – 2.58 |
| | 2004 | 66 | 1.73 | 1.34 – 2.20 |
| | 2005 | 79 | 2.05 | 1.62 – 2.55 |
| | 2006 | 81 | 2.03 | 1.61 – 2.52 |
| | 2007 | 88 | 2.16 | 1.73 – 2.66 |
| | 2008 | 78 | 1.92 | 1.52 – 2.40 |
| | 2009 | 89 | 2.22 | 1.78 – 2.73 |
| | 2010 | 76 | 1.97 | 1.55 – 2.47 |
| | 2011 | 89 | 2.36 | 1.89 – 2.90 |
| | 2012 | 93 | 2.43 | 1.96 – 2.98 |
| | 2013 | 99 | 2.56 | 2.08 – 3.11 |
| | 2014 | 90 | 2.25 | 1.81 – 2.77 |
| | 2015 | 89 | 2.21 | 1.77 – 2.71 |
| | 2016 | 97 | 2.44 | 1.98 – 2.98 |
| 2017 | 89 | 2.33 | 1.87 – 2.87 | |
| 2018 | 78 | 2.07 | 1.64 – 2.59 | |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|--|------|---------------|-------------------|--|
| | 2019 | 86 | 2.28 | 1.82 – 2.81 |
| | 2020 | 93 | 2.52 | 2.04 – 3.09 |
| Pyloric stenosis | 1999 | 644 | 18.44 | 17.02 – 19.87 |
| | 2000 | 752 | 20.70 | 19.22 – 22.18 |
| | 2001 | 719 | 19.69 | 18.25 – 21.13 |
| | 2002 | 645 | 17.32 | 15.98 – 18.66 |
| | 2003 | 585 | 15.50 | 14.25 – 16.76 |
| | 2004 | 777 | 20.37 | 18.94 – 21.80 |
| | 2005 | 693 | 17.97 | 16.64 – 19.31 |
| | 2006 | 980 | 24.54 | 23.01 – 26.08 |
| | 2007 | 917 | 22.51 | 21.05 – 23.96 |
| | 2008 | 715 | 17.64 | 16.35 – 18.94 |
| | 2009 | 630 | 15.69 | 14.46 – 16.91 |
| | 2010 | 576 | 14.93 | 13.71 – 16.15 |
| | 2011 | 543 | 14.39 | 13.18 – 15.60 |
| | 2012 | 449 | 11.74 | 10.65 – 12.83 |
| | 2013 | 460 | 11.88 | 10.80 – 12.97 |
| | 2014 | 538 | 13.47 | 12.33 – 14.61 |
| | 2015 | 499 | 12.37 | 11.28 – 13.45 |
| | 2016 | 543 | 13.68 | 12.53 – 14.83 |
| | 2017 | 422 | 11.05 | 10.00 – 12.11 |
| | 2018 | 553 | 14.69 | 13.46 – 15.91 |
| | 2019 | 501 | 13.26 | 12.10 – 14.43 |
| | 2020 | 447 | 12.14 | 11.01 – 13.26 |
| Stenosis or atresia of the small intestine | 1999 | 92 | 2.63 | 2.12 – 3.23 |
| | 2000 | 114 | 3.14 | 2.56 – 3.71 |
| | 2001 | 130 | 3.56 | 2.95 – 4.17 |
| | 2002 | 111 | 2.98 | 2.43 – 3.54 |
| | 2003 | 96 | 2.54 | 2.06 – 3.11 |
| | 2004 | 126 | 3.30 | 2.73 – 3.88 |
| | 2005 | 113 | 2.93 | 2.39 – 3.47 |
| | 2006 | 132 | 3.31 | 2.74 – 3.87 |
| | 2007 | 152 | 3.73 | 3.14 – 4.32 |
| | 2008 | 147 | 3.63 | 3.04 – 4.21 |
| | 2009 | 139 | 3.46 | 2.89 – 4.04 |
| | 2010 | 146 | 3.78 | 3.17 – 4.40 |
| | 2011 | 129 | 3.42 | 2.83 – 4.01 |
| | 2012 | 125 | 3.27 | 2.70 – 3.84 |
| | 2013 | 138 | 3.56 | 2.97 – 4.16 |
| | 2014 | 145 | 3.63 | 3.04 – 4.22 |
| | 2015 | 149 | 3.69 | 3.10 – 4.29 |
| | 2016 | 129 | 3.25 | 2.69 – 3.81 |
| | 2017 | 150 | 3.93 | 3.30 – 4.56 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|---|------|---------------|-------------------|--|
| | 2018 | 151 | 4.01 | 3.37 – 4.65 |
| | 2019 | 142 | 3.76 | 3.14 – 4.38 |
| | 2020 | 119 | 3.23 | 2.65 – 3.81 |
| Stenosis or atresia of large intestine, rectum, or anal canal | 1999 | 192 | 5.50 | 4.72 – 6.28 |
| | 2000 | 164 | 4.51 | 3.82 – 5.20 |
| | 2001 | 191 | 5.23 | 4.49 – 5.97 |
| | 2002 | 201 | 5.40 | 4.65 – 6.14 |
| | 2003 | 212 | 5.62 | 4.86 – 6.37 |
| | 2004 | 192 | 5.03 | 4.32 – 5.75 |
| | 2005 | 213 | 5.52 | 4.78 – 6.27 |
| | 2006 | 210 | 5.26 | 4.55 – 5.97 |
| | 2007 | 213 | 5.23 | 4.53 – 5.93 |
| | 2008 | 221 | 5.45 | 4.73 – 6.17 |
| | 2009 | 222 | 5.53 | 4.80 – 6.26 |
| | 2010 | 201 | 5.21 | 4.49 – 5.93 |
| | 2011 | 209 | 5.54 | 4.79 – 6.29 |
| | 2012 | 201 | 5.26 | 4.53 – 5.98 |
| | 2013 | 210 | 5.42 | 4.69 – 6.16 |
| | 2014 | 218 | 5.46 | 4.73 – 6.18 |
| | 2015 | 199 | 4.93 | 4.25 – 5.62 |
| | 2016 | 211 | 5.31 | 4.60 – 6.03 |
| | 2017 | 208 | 5.45 | 4.71 – 6.19 |
| | 2018 | 213 | 5.66 | 4.90 – 6.42 |
| | 2019 | 200 | 5.30 | 4.56 – 6.03 |
| | 2020 | 194 | 5.27 | 4.53 – 6.01 |
| Hirschsprung disease | 1999 | 39 | 1.12 | 0.79 – 1.53 |
| | 2000 | 47 | 1.29 | 0.95 – 1.72 |
| | 2001 | 42 | 1.15 | 0.83 – 1.55 |
| | 2002 | 44 | 1.18 | 0.86 – 1.59 |
| | 2003 | 44 | 1.17 | 0.85 – 1.57 |
| | 2004 | 52 | 1.36 | 1.02 – 1.79 |
| | 2005 | 57 | 1.48 | 1.12 – 1.92 |
| | 2006 | 48 | 1.20 | 0.89 – 1.59 |
| | 2007 | 67 | 1.64 | 1.27 – 2.09 |
| | 2008 | 63 | 1.55 | 1.19 – 1.99 |
| | 2009 | 50 | 1.25 | 0.92 – 1.64 |
| | 2010 | 56 | 1.45 | 1.10 – 1.89 |
| | 2011 | 62 | 1.64 | 1.26 – 2.11 |
| | 2012 | 55 | 1.44 | 1.08 – 1.87 |
| | 2013 | 59 | 1.52 | 1.16 – 1.97 |
| | 2014 | 57 | 1.43 | 1.08 – 1.85 |
| | 2015 | 74 | 1.83 | 1.44 – 2.30 |
| | 2016 | 71 | 1.79 | 1.40 – 2.26 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|----------------------------------|------|---------------|-------------------|--|
| | 2017 | 51 | 1.34 | 0.99 – 1.76 |
| | 2018 | 62 | 1.65 | 1.26 – 2.11 |
| | 2019 | 58 | 1.54 | 1.17 – 1.99 |
| | 2020 | 55 | 1.49 | 1.12 – 1.94 |
| Biliary atresia | 1999 | 27 | 0.77 | 0.51 – 1.13 |
| | 2000 | 28 | 0.77 | 0.51 – 1.11 |
| | 2001 | 24 | 0.66 | 0.42 – 0.98 |
| | 2002 | 24 | 0.64 | 0.41 – 0.96 |
| | 2003 | 31 | 0.82 | 0.56 – 1.17 |
| | 2004 | 28 | 0.73 | 0.49 – 1.06 |
| | 2005 | 26 | 0.67 | 0.44 – 0.99 |
| | 2006 | 33 | 0.83 | 0.57 – 1.16 |
| | 2007 | 30 | 0.74 | 0.50 – 1.05 |
| | 2008 | 26 | 0.64 | 0.42 – 0.94 |
| | 2009 | 33 | 0.82 | 0.57 – 1.15 |
| | 2010 | 27 | 0.70 | 0.46 – 1.02 |
| | 2011 | 30 | 0.80 | 0.54 – 1.14 |
| | 2012 | 25 | 0.65 | 0.42 – 0.96 |
| | 2013 | 23 | 0.59 | 0.38 – 0.89 |
| | 2014 | 21 | 0.53 | 0.33 – 0.80 |
| | 2015 | 33 | 0.82 | 0.56 – 1.15 |
| | 2016 | 29 | 0.73 | 0.49 – 1.05 |
| | 2017 | 31 | 0.81 | 0.55 – 1.15 |
| | 2018 | 22 | 0.58 | 0.37 – 0.88 |
| | 2019 | 24 | 0.64 | 0.41 – 0.95 |
| | 2020 | 28 | 0.76 | 0.51 – 1.10 |
| Genitourinary | | | | |
| Hypospadias (among males) | 1999 | 1,018 | 57.05 | 53.54 – 60.55 |
| | 2000 | 1,032 | 55.61 | 52.21 – 59.00 |
| | 2001 | 979 | 52.42 | 49.13 – 55.70 |
| | 2002 | 987 | 51.90 | 48.67 – 55.14 |
| | 2003 | 1,009 | 52.39 | 49.16 – 55.63 |
| | 2004 | 1,041 | 53.38 | 50.14 – 56.62 |
| | 2005 | 1,131 | 57.27 | 53.93 – 60.61 |
| | 2006 | 1,102 | 54.01 | 50.82 – 57.20 |
| | 2007 | 1,147 | 55.09 | 51.90 – 58.27 |
| | 2008 | 1,210 | 58.31 | 55.03 – 61.60 |
| | 2009 | 1,271 | 62.04 | 58.63 – 65.45 |
| | 2010 | 1,259 | 63.94 | 60.41 – 67.47 |
| | 2011 | 1,230 | 63.81 | 60.24 – 67.37 |
| | 2012 | 1,238 | 63.32 | 59.79 – 66.84 |
| | 2013 | 1,362 | 68.75 | 65.10 – 72.40 |
| | 2014 | 1,484 | 72.57 | 68.87 – 76.26 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|------------------------------|------|---------------|-------------------|--|
| | 2015 | 1,400 | 67.97 | 64.41 – 71.53 |
| | 2016 | 1,511 | 74.50 | 70.75 – 78.26 |
| | 2017 | 1,421 | 72.92 | 69.13 – 76.71 |
| | 2018 | 1,461 | 76.00 | 72.10 – 79.90 |
| | 2019 | 1,267 | 65.64 | 62.02 – 69.25 |
| | 2020 | 1,239 | 65.71 | 62.05 – 69.37 |
| Epispadias | 1999 | 21 | 0.60 | 0.37 – 0.92 |
| | 2000 | 32 | 0.88 | 0.60 – 1.24 |
| | 2001 | 29 | 0.79 | 0.53 – 1.14 |
| | 2002 | 22 | 0.59 | 0.37 – 0.89 |
| | 2003 | 35 | 0.93 | 0.65 – 1.29 |
| | 2004 | 31 | 0.81 | 0.55 – 1.15 |
| | 2005 | 27 | 0.70 | 0.46 – 1.02 |
| | 2006 | 42 | 1.05 | 0.76 – 1.42 |
| | 2007 | 39 | 0.96 | 0.68 – 1.31 |
| | 2008 | 40 | 0.99 | 0.71 – 1.34 |
| | 2009 | 41 | 1.02 | 0.73 – 1.38 |
| | 2010 | 44 | 1.14 | 0.83 – 1.53 |
| | 2011 | 39 | 1.03 | 0.74 – 1.41 |
| | 2012 | 55 | 1.44 | 1.08 – 1.87 |
| | 2013 | 80 | 2.07 | 1.64 – 2.57 |
| | 2014 | 50 | 1.25 | 0.93 – 1.65 |
| | 2015 | 61 | 1.51 | 1.16 – 1.94 |
| | 2016 | 49 | 1.23 | 0.91 – 1.63 |
| | 2017 | 40 | 1.05 | 0.75 – 1.43 |
| | 2018 | 23 | 0.61 | 0.39 – 0.92 |
| | 2019 | 14 | 0.37 | 0.20 – 0.62 |
| | 2020 | 6 | 0.16 | 0.06 – 0.35 |
| Renal agenesis or dysgenesis | 1999 | 200 | 5.73 | 4.93 – 6.52 |
| | 2000 | 177 | 4.87 | 4.15 – 5.59 |
| | 2001 | 193 | 5.29 | 4.54 – 6.03 |
| | 2002 | 181 | 4.86 | 4.15 – 5.57 |
| | 2003 | 222 | 5.88 | 5.11 – 6.66 |
| | 2004 | 212 | 5.56 | 4.81 – 6.31 |
| | 2005 | 189 | 4.90 | 4.20 – 5.60 |
| | 2006 | 240 | 6.01 | 5.25 – 6.77 |
| | 2007 | 247 | 6.06 | 5.31 – 6.82 |
| | 2008 | 241 | 5.95 | 5.20 – 6.70 |
| | 2009 | 260 | 6.47 | 5.69 – 7.26 |
| | 2010 | 263 | 6.82 | 5.99 – 7.64 |
| | 2011 | 262 | 6.94 | 6.10 – 7.79 |
| | 2012 | 241 | 6.30 | 5.51 – 7.10 |
| | 2013 | 286 | 7.39 | 6.53 – 8.24 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|--|------|---------------|-------------------|--|
| | 2014 | 264 | 6.61 | 5.81 – 7.41 |
| | 2015 | 256 | 6.35 | 5.57 – 7.12 |
| | 2016 | 283 | 7.13 | 6.30 – 7.96 |
| | 2017 | 292 | 7.65 | 6.77 – 8.52 |
| | 2018 | 313 | 8.31 | 7.39 – 9.23 |
| | 2019 | 348 | 9.21 | 8.25 – 10.18 |
| | 2020 | 298 | 8.09 | 7.17 – 9.01 |
| Bladder exstrophy | 1999 | 4 | 0.11 | 0.03 – 0.29 |
| | 2000 | 11 | 0.30 | 0.15 – 0.54 |
| | 2001 | 5 | 0.14 | 0.04 – 0.32 |
| | 2002 | 12 | 0.32 | 0.17 – 0.56 |
| | 2003 | 11 | 0.29 | 0.15 – 0.52 |
| | 2004 | 10 | 0.26 | 0.13 – 0.48 |
| | 2005 | 3 | 0.08 | 0.02 – 0.23 |
| | 2006 | 6 | 0.15 | 0.06 – 0.33 |
| | 2007 | 7 | 0.17 | 0.07 – 0.35 |
| | 2008 | 8 | 0.20 | 0.09 – 0.39 |
| | 2009 | 5 | 0.12 | 0.04 – 0.29 |
| | 2010 | 9 | 0.23 | 0.11 – 0.44 |
| | 2011 | 6 | 0.16 | 0.06 – 0.35 |
| | 2012 | 7 | 0.18 | 0.07 – 0.38 |
| | 2013 | 11 | 0.28 | 0.14 – 0.51 |
| | 2014 | 5 | 0.13 | 0.04 – 0.29 |
| | 2015 | 7 | 0.17 | 0.07 – 0.36 |
| | 2016 | 2 | 0.05 | 0.01 – 0.18 |
| | 2017 | 3 | 0.08 | 0.02 – 0.23 |
| | 2018 | 3 | 0.08 | 0.02 – 0.23 |
| | 2019 | 9 | 0.24 | 0.11 – 0.45 |
| | 2020 | 4 | 0.11 | 0.03 – 0.28 |
| Musculoskeletal | | | | |
| Congenital hip dislocation without hip dysplasia | 1999 | 169 | 4.84 | 4.11 – 5.57 |
| | 2000 | 149 | 4.10 | 3.44 – 4.76 |
| | 2001 | 127 | 3.48 | 2.87 – 4.08 |
| | 2002 | 110 | 2.95 | 2.40 – 3.51 |
| | 2003 | 124 | 3.29 | 2.71 – 3.86 |
| | 2004 | 113 | 2.96 | 2.42 – 3.51 |
| | 2005 | 107 | 2.78 | 2.25 – 3.30 |
| | 2006 | 107 | 2.68 | 2.17 – 3.19 |
| | 2007 | 115 | 2.82 | 2.31 – 3.34 |
| | 2008 | 96 | 2.37 | 1.92 – 2.89 |
| | 2009 | 108 | 2.69 | 2.18 – 3.20 |
| | 2010 | 111 | 2.88 | 2.34 – 3.41 |
| | 2011 | 76 | 2.01 | 1.59 – 2.52 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|--------------------------------------|------|---------------|-------------------|--|
| | 2012 | 71 | 1.86 | 1.45 – 2.34 |
| | 2013 | 88 | 2.27 | 1.82 – 2.80 |
| | 2014 | 74 | 1.85 | 1.45 – 2.33 |
| | 2015 | 80 | 1.98 | 1.57 – 2.47 |
| | 2016 | 75 | 1.89 | 1.49 – 2.37 |
| | 2017 | 76 | 1.99 | 1.57 – 2.49 |
| | 2018 | 54 | 1.43 | 1.08 – 1.87 |
| | 2019 | 47 | 1.24 | 0.91 – 1.65 |
| | 2020 | 45 | 1.22 | 0.89 – 1.63 |
| Talipes equinovarus / clubfoot | 1999 | 500 | 14.32 | 13.06 – 15.58 |
| | 2000 | 519 | 14.28 | 13.06 – 15.51 |
| | 2001 | 536 | 14.68 | 13.44 – 15.92 |
| | 2002 | 572 | 15.36 | 14.10 – 16.62 |
| | 2003 | 574 | 15.21 | 13.97 – 16.45 |
| | 2004 | 583 | 15.28 | 14.04 – 16.52 |
| | 2005 | 553 | 14.34 | 13.15 – 15.54 |
| | 2006 | 580 | 14.53 | 13.34 – 15.71 |
| | 2007 | 626 | 15.36 | 14.16 – 16.57 |
| | 2008 | 620 | 15.30 | 14.10 – 16.50 |
| | 2009 | 646 | 16.09 | 14.85 – 17.33 |
| | 2010 | 645 | 16.72 | 15.43 – 18.01 |
| | 2011 | 643 | 17.04 | 15.73 – 18.36 |
| | 2012 | 653 | 17.07 | 15.77 – 18.38 |
| | 2013 | 646 | 16.69 | 15.40 – 17.97 |
| | 2014 | 736 | 18.42 | 17.09 – 19.75 |
| | 2015 | 699 | 17.33 | 16.04 – 18.61 |
| | 2016 | 766 | 19.29 | 17.93 – 20.66 |
| | 2017 | 749 | 19.61 | 18.21 – 21.02 |
| | 2018 | 743 | 19.73 | 18.32 – 21.15 |
| | 2019 | 730 | 19.33 | 17.92 – 20.73 |
| | 2020 | 695 | 18.87 | 17.47 – 20.27 |
| Reduction defects of the upper limbs | 1999 | 141 | 4.04 | 3.37 – 4.70 |
| | 2000 | 157 | 4.32 | 3.65 – 5.00 |
| | 2001 | 148 | 4.05 | 3.40 – 4.71 |
| | 2002 | 148 | 3.97 | 3.33 – 4.61 |
| | 2003 | 151 | 4.00 | 3.36 – 4.64 |
| | 2004 | 149 | 3.91 | 3.28 – 4.53 |
| | 2005 | 156 | 4.05 | 3.41 – 4.68 |
| | 2006 | 164 | 4.11 | 3.48 – 4.74 |
| | 2007 | 167 | 4.10 | 3.48 – 4.72 |
| | 2008 | 169 | 4.17 | 3.54 – 4.80 |
| | 2009 | 168 | 4.18 | 3.55 – 4.82 |
| | 2010 | 194 | 5.03 | 4.32 – 5.74 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|--------------------------------------|------|---------------|-------------------|--|
| | 2011 | 183 | 4.85 | 4.15 – 5.55 |
| | 2012 | 140 | 3.66 | 3.05 – 4.27 |
| | 2013 | 161 | 4.16 | 3.52 – 4.80 |
| | 2014 | 153 | 3.83 | 3.22 – 4.44 |
| | 2015 | 176 | 4.36 | 3.72 – 5.01 |
| | 2016 | 164 | 4.13 | 3.50 – 4.76 |
| | 2017 | 146 | 3.82 | 3.20 – 4.44 |
| | 2018 | 161 | 4.28 | 3.62 – 4.94 |
| | 2019 | 158 | 4.18 | 3.53 – 4.84 |
| | 2020 | 147 | 3.99 | 3.35 – 4.64 |
| Reduction defects of the lower limbs | 1999 | 81 | 2.32 | 1.84 – 2.88 |
| | 2000 | 55 | 1.51 | 1.14 – 1.97 |
| | 2001 | 79 | 2.16 | 1.71 – 2.70 |
| | 2002 | 63 | 1.69 | 1.30 – 2.16 |
| | 2003 | 75 | 1.99 | 1.56 – 2.49 |
| | 2004 | 64 | 1.68 | 1.29 – 2.14 |
| | 2005 | 83 | 2.15 | 1.71 – 2.67 |
| | 2006 | 72 | 1.80 | 1.41 – 2.27 |
| | 2007 | 89 | 2.18 | 1.75 – 2.69 |
| | 2008 | 83 | 2.05 | 1.63 – 2.54 |
| | 2009 | 75 | 1.87 | 1.47 – 2.34 |
| | 2010 | 84 | 2.18 | 1.74 – 2.70 |
| | 2011 | 79 | 2.09 | 1.66 – 2.61 |
| | 2012 | 84 | 2.20 | 1.75 – 2.72 |
| | 2013 | 65 | 1.68 | 1.30 – 2.14 |
| | 2014 | 81 | 2.03 | 1.61 – 2.52 |
| | 2015 | 82 | 2.03 | 1.62 – 2.52 |
| | 2016 | 69 | 1.74 | 1.35 – 2.20 |
| | 2017 | 78 | 2.04 | 1.61 – 2.55 |
| | 2018 | 56 | 1.49 | 1.12 – 1.93 |
| | 2019 | 86 | 2.28 | 1.82 – 2.81 |
| | 2020 | 75 | 2.04 | 1.60 – 2.55 |
| Craniosynostosis | 1999 | 141 | 4.04 | 3.37 – 4.70 |
| | 2000 | 165 | 4.54 | 3.85 – 5.23 |
| | 2001 | 139 | 3.81 | 3.17 – 4.44 |
| | 2002 | 173 | 4.65 | 3.95 – 5.34 |
| | 2003 | 168 | 4.45 | 3.78 – 5.13 |
| | 2004 | 172 | 4.51 | 3.84 – 5.18 |
| | 2005 | 159 | 4.12 | 3.48 – 4.77 |
| | 2006 | 180 | 4.51 | 3.85 – 5.17 |
| | 2007 | 217 | 5.33 | 4.62 – 6.03 |
| | 2008 | 213 | 5.26 | 4.55 – 5.96 |
| | 2009 | 208 | 5.18 | 4.48 – 5.88 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|----------------------------|------|------------------|----------------------|---|
| | 2010 | 258 | 6.69 | 5.87 – 7.50 |
| | 2011 | 231 | 6.12 | 5.33 – 6.91 |
| | 2012 | 273 | 7.14 | 6.29 – 7.99 |
| | 2013 | 226 | 5.84 | 5.08 – 6.60 |
| | 2014 | 248 | 6.21 | 5.44 – 6.98 |
| | 2015 | 254 | 6.30 | 5.52 – 7.07 |
| | 2016 | 236 | 5.94 | 5.19 – 6.70 |
| | 2017 | 250 | 6.55 | 5.74 – 7.36 |
| | 2018 | 285 | 7.57 | 6.69 – 8.45 |
| | 2019 | 270 | 7.15 | 6.30 – 8.00 |
| | 2020 | 250 | 6.79 | 5.95 – 7.63 |
| Achondroplasia | 1999 | 12 | 0.34 | 0.18 – 0.60 |
| | 2000 | 10 | 0.28 | 0.13 – 0.51 |
| | 2001 | 13 | 0.36 | 0.19 – 0.61 |
| | 2002 | 10 | 0.27 | 0.13 – 0.49 |
| | 2003 | 11 | 0.29 | 0.15 – 0.52 |
| | 2004 | 9 | 0.24 | 0.11 – 0.45 |
| | 2005 | 10 | 0.26 | 0.12 – 0.48 |
| | 2006 | 9 | 0.23 | 0.10 – 0.43 |
| | 2007 | 19 | 0.47 | 0.28 – 0.73 |
| | 2008 | 12 | 0.30 | 0.15 – 0.52 |
| | 2009 | 16 | 0.40 | 0.23 – 0.65 |
| | 2010 | 13 | 0.34 | 0.18 – 0.58 |
| | 2011 | 11 | 0.29 | 0.15 – 0.52 |
| | 2012 | 15 | 0.39 | 0.22 – 0.65 |
| | 2013 | 16 | 0.41 | 0.24 – 0.67 |
| | 2014 | 13 | 0.33 | 0.17 – 0.56 |
| | 2015 | 17 | 0.42 | 0.25 – 0.67 |
| | 2016 | 23 | 0.58 | 0.37 – 0.87 |
| | 2017 | 14 | 0.37 | 0.20 – 0.62 |
| | 2018 | 24 | 0.64 | 0.41 – 0.95 |
| | 2019 | 13 | 0.34 | 0.18 – 0.59 |
| | 2020 | 17 | 0.46 | 0.27 – 0.74 |
| Diaphragmatic hernia | 1999 | 112 | 3.21 | 2.61 – 3.80 |
| | 2000 | 99 | 2.72 | 2.21 – 3.32 |
| | 2001 | 83 | 2.27 | 1.81 – 2.82 |
| | 2002 | 95 | 2.55 | 2.06 – 3.12 |
| | 2003 | 108 | 2.86 | 2.32 – 3.40 |
| | 2004 | 100 | 2.62 | 2.13 – 3.19 |
| | 2005 | 97 | 2.52 | 2.04 – 3.07 |
| | 2006 | 116 | 2.91 | 2.38 – 3.43 |
| | 2007 | 128 | 3.14 | 2.60 – 3.69 |
| | 2008 | 115 | 2.84 | 2.32 – 3.36 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|----------------------------|------|------------------|----------------------|---|
| | 2009 | 107 | 2.66 | 2.16 – 3.17 |
| | 2010 | 121 | 3.14 | 2.58 – 3.70 |
| | 2011 | 100 | 2.65 | 2.16 – 3.22 |
| | 2012 | 107 | 2.80 | 2.27 – 3.33 |
| | 2013 | 121 | 3.13 | 2.57 – 3.68 |
| | 2014 | 110 | 2.75 | 2.24 – 3.27 |
| | 2015 | 97 | 2.40 | 1.95 – 2.93 |
| | 2016 | 105 | 2.64 | 2.14 – 3.15 |
| | 2017 | 121 | 3.17 | 2.60 – 3.73 |
| | 2018 | 107 | 2.84 | 2.30 – 3.38 |
| | 2019 | 93 | 2.46 | 1.99 – 3.02 |
| | 2020 | 117 | 3.18 | 2.60 – 3.75 |
| Omphalocele | 1999 | 78 | 2.23 | 1.77 – 2.79 |
| | 2000 | 83 | 2.28 | 1.82 – 2.83 |
| | 2001 | 85 | 2.33 | 1.86 – 2.88 |
| | 2002 | 77 | 2.07 | 1.63 – 2.58 |
| | 2003 | 73 | 1.93 | 1.52 – 2.43 |
| | 2004 | 83 | 2.18 | 1.73 – 2.70 |
| | 2005 | 73 | 1.89 | 1.48 – 2.38 |
| | 2006 | 86 | 2.15 | 1.72 – 2.66 |
| | 2007 | 80 | 1.96 | 1.56 – 2.44 |
| | 2008 | 79 | 1.95 | 1.54 – 2.43 |
| | 2009 | 90 | 2.24 | 1.80 – 2.75 |
| | 2010 | 79 | 2.05 | 1.62 – 2.55 |
| | 2011 | 69 | 1.83 | 1.42 – 2.31 |
| | 2012 | 87 | 2.27 | 1.82 – 2.81 |
| | 2013 | 85 | 2.20 | 1.75 – 2.72 |
| | 2014 | 86 | 2.15 | 1.72 – 2.66 |
| | 2015 | 96 | 2.38 | 1.93 – 2.91 |
| | 2016 | 86 | 2.17 | 1.73 – 2.68 |
| | 2017 | 94 | 2.46 | 1.99 – 3.01 |
| | 2018 | 84 | 2.23 | 1.78 – 2.76 |
| | 2019 | 82 | 2.17 | 1.73 – 2.69 |
| | 2020 | 86 | 2.33 | 1.87 – 2.88 |
| Gastroschisis | 1999 | 137 | 3.92 | 3.27 – 4.58 |
| | 2000 | 143 | 3.94 | 3.29 – 4.58 |
| | 2001 | 148 | 4.05 | 3.40 – 4.71 |
| | 2002 | 149 | 4.00 | 3.36 – 4.64 |
| | 2003 | 173 | 4.58 | 3.90 – 5.27 |
| | 2004 | 165 | 4.33 | 3.67 – 4.99 |
| | 2005 | 208 | 5.40 | 4.66 – 6.13 |
| | 2006 | 204 | 5.11 | 4.41 – 5.81 |
| | 2007 | 257 | 6.31 | 5.54 – 7.08 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|-----------------------------|------|---------------|-------------------|--|
| | 2008 | 250 | 6.17 | 5.40 – 6.93 |
| | 2009 | 242 | 6.03 | 5.27 – 6.79 |
| | 2010 | 255 | 6.61 | 5.80 – 7.42 |
| | 2011 | 216 | 5.73 | 4.96 – 6.49 |
| | 2012 | 224 | 5.86 | 5.09 – 6.62 |
| | 2013 | 220 | 5.68 | 4.93 – 6.43 |
| | 2014 | 232 | 5.81 | 5.06 – 6.55 |
| | 2015 | 231 | 5.73 | 4.99 – 6.46 |
| | 2016 | 188 | 4.74 | 4.06 – 5.41 |
| | 2017 | 176 | 4.61 | 3.93 – 5.29 |
| | 2018 | 182 | 4.83 | 4.13 – 5.54 |
| | 2019 | 172 | 4.55 | 3.87 – 5.23 |
| | 2020 | 144 | 3.91 | 3.27 – 4.55 |
| Chromosomal | | | | |
| Trisomy 21 (Down syndrome) | 1999 | 426 | 12.20 | 11.04 – 13.36 |
| | 2000 | 476 | 13.10 | 11.92 – 14.28 |
| | 2001 | 494 | 13.53 | 12.34 – 14.72 |
| | 2002 | 455 | 12.22 | 11.10 – 13.34 |
| | 2003 | 457 | 12.11 | 11.00 – 13.22 |
| | 2004 | 513 | 13.45 | 12.29 – 14.61 |
| | 2005 | 482 | 12.50 | 11.39 – 13.62 |
| | 2006 | 559 | 14.00 | 12.84 – 15.16 |
| | 2007 | 550 | 13.50 | 12.37 – 14.63 |
| | 2008 | 544 | 13.42 | 12.30 – 14.55 |
| | 2009 | 556 | 13.84 | 12.69 – 15.00 |
| | 2010 | 580 | 15.04 | 13.81 – 16.26 |
| | 2011 | 553 | 14.66 | 13.44 – 15.88 |
| | 2012 | 571 | 14.93 | 13.71 – 16.16 |
| | 2013 | 551 | 14.23 | 13.05 – 15.42 |
| | 2014 | 561 | 14.04 | 12.88 – 15.21 |
| | 2015 | 524 | 12.99 | 11.88 – 14.10 |
| | 2016 | 574 | 14.46 | 13.28 – 15.64 |
| | 2017 | 553 | 14.48 | 13.27 – 15.69 |
| | 2018 | 587 | 15.59 | 14.33 – 16.85 |
| | 2019 | 576 | 15.25 | 14.00 – 16.50 |
| | 2020 | 552 | 14.99 | 13.74 – 16.24 |
| Trisomy 13 (Patau syndrome) | 1999 | 43 | 1.23 | 0.89 – 1.66 |
| | 2000 | 51 | 1.40 | 1.05 – 1.85 |
| | 2001 | 42 | 1.15 | 0.83 – 1.55 |
| | 2002 | 43 | 1.15 | 0.84 – 1.56 |
| | 2003 | 42 | 1.11 | 0.80 – 1.50 |
| | 2004 | 46 | 1.21 | 0.88 – 1.61 |
| | 2005 | 34 | 0.88 | 0.61 – 1.23 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|--|------|------------------|----------------------|---|
| | 2006 | 42 | 1.05 | 0.76 – 1.42 |
| | 2007 | 51 | 1.25 | 0.93 – 1.65 |
| | 2008 | 45 | 1.11 | 0.81 – 1.49 |
| | 2009 | 53 | 1.32 | 0.99 – 1.73 |
| | 2010 | 59 | 1.53 | 1.16 – 1.97 |
| | 2011 | 43 | 1.14 | 0.82 – 1.54 |
| | 2012 | 53 | 1.39 | 1.04 – 1.81 |
| | 2013 | 43 | 1.11 | 0.80 – 1.50 |
| | 2014 | 45 | 1.13 | 0.82 – 1.51 |
| | 2015 | 38 | 0.94 | 0.67 – 1.29 |
| | 2016 | 51 | 1.28 | 0.96 – 1.69 |
| | 2017 | 41 | 1.07 | 0.77 – 1.46 |
| | 2018 | 39 | 1.04 | 0.74 – 1.42 |
| | 2019 | 28 | 0.74 | 0.49 – 1.07 |
| | 2020 | 38 | 1.03 | 0.73 – 1.42 |
| <hr/> | | | | |
| Trisomy 18 (Edwards syndrome) | 1999 | 84 | 2.41 | 1.92 – 2.98 |
| | 2000 | 78 | 2.15 | 1.70 – 2.68 |
| | 2001 | 81 | 2.22 | 1.76 – 2.76 |
| | 2002 | 82 | 2.20 | 1.75 – 2.73 |
| | 2003 | 80 | 2.12 | 1.68 – 2.64 |
| | 2004 | 97 | 2.54 | 2.06 – 3.10 |
| | 2005 | 96 | 2.49 | 2.02 – 3.04 |
| | 2006 | 110 | 2.75 | 2.24 – 3.27 |
| | 2007 | 102 | 2.50 | 2.02 – 2.99 |
| | 2008 | 120 | 2.96 | 2.43 – 3.49 |
| | 2009 | 120 | 2.99 | 2.45 – 3.52 |
| | 2010 | 99 | 2.57 | 2.09 – 3.12 |
| | 2011 | 109 | 2.89 | 2.35 – 3.43 |
| | 2012 | 112 | 2.93 | 2.39 – 3.47 |
| | 2013 | 95 | 2.45 | 1.99 – 3.00 |
| | 2014 | 116 | 2.90 | 2.38 – 3.43 |
| | 2015 | 87 | 2.16 | 1.73 – 2.66 |
| | 2016 | 107 | 2.70 | 2.18 – 3.21 |
| | 2017 | 84 | 2.20 | 1.75 – 2.72 |
| | 2018 | 86 | 2.28 | 1.83 – 2.82 |
| | 2019 | 90 | 2.38 | 1.92 – 2.93 |
| | 2020 | 90 | 2.44 | 1.96 – 3.00 |
| <hr/> | | | | |
| Infants and fetuses with regular reportable birth defects | 1999 | 11,501 | 329.39 | 323.37 – 335.41 |
| | 2000 | 12,445 | 342.53 | 336.51 – 348.55 |
| | 2001 | 12,382 | 339.15 | 333.17 – 345.12 |
| | 2002 | 13,116 | 352.23 | 346.20 – 358.26 |
| | 2003 | 13,867 | 367.46 | 361.34 – 373.58 |
| | 2004 | 15,195 | 398.36 | 392.02 – 404.69 |

| Birth Defect (Body System) | Year | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|----------------------------|------|---------------|-------------------|--|
| | 2005 | 15,992 | 414.80 | 408.37 – 421.23 |
| | 2006 | 16,691 | 418.00 | 411.66 – 424.34 |
| | 2007 | 17,448 | 428.22 | 421.87 – 434.58 |
| | 2008 | 18,271 | 450.87 | 444.33 – 457.40 |
| | 2009 | 19,091 | 475.37 | 468.63 – 482.12 |
| | 2010 | 19,686 | 510.34 | 503.21 – 517.46 |
| | 2011 | 19,769 | 524.00 | 516.69 – 531.30 |
| | 2012 | 20,330 | 531.59 | 524.28 – 538.90 |
| | 2013 | 21,471 | 554.65 | 547.23 – 562.07 |
| | 2014 | 22,595 | 565.61 | 558.23 – 572.98 |
| | 2015 | 23,275 | 576.91 | 569.50 – 584.33 |
| | 2016 | 25,657 | 646.27 | 638.37 – 654.18 |
| | 2017 | 24,281 | 635.83 | 627.84 – 643.83 |
| | 2018 | 25,494 | 677.12 | 668.81 – 685.43 |
| | 2019 | 25,259 | 668.74 | 660.49 – 676.99 |
| | 2020 | 25,917 | 703.66 | 695.09 – 712.23 |

Prevalence (rate) is expressed as the number of cases per 10,000 live births.

Please see the Methods section of the Annual Report for additional information:

<https://www.dshs.texas.gov/sites/default/files/birthdefects/annualreport/1999-2020-TBDR-Methods.pdf>

Prepared by: Texas Birth Defects Registry, Birth Defects Epidemiology and Surveillance Branch, Texas Department of State Health Services, February 2024.

Texas Birth Defects Registry (TBDR) Annual Report

Table 2B. Prevalence of Selected Birth Defects by 5-Year Period, Texas, 2001–2020

| Birth Defect (Body System) | 5-Year Period | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|---|---------------|------------------|----------------------|---|
| Central Nervous System | | | | |
| Anencephaly | 2001-2005 | 462 | 2.46 | 2.23 – 2.68 |
| | 2006-2010 | 519 | 2.60 | 2.37 – 2.82 |
| | 2011-2015 | 461 | 2.36 | 2.15 – 2.58 |
| | 2016-2020 | 441 | 2.32 | 2.10 – 2.54 |
| Spina bifida without anencephaly | 2001-2005 | 672 | 3.57 | 3.30 – 3.84 |
| | 2006-2010 | 745 | 3.73 | 3.46 – 3.99 |
| | 2011-2015 | 750 | 3.85 | 3.57 – 4.12 |
| | 2016-2020 | 686 | 3.61 | 3.34 – 3.88 |
| Encephalocele | 2001-2005 | 150 | 0.80 | 0.67 – 0.92 |
| | 2006-2010 | 201 | 1.01 | 0.87 – 1.14 |
| | 2011-2015 | 192 | 0.98 | 0.85 – 1.12 |
| | 2016-2020 | 199 | 1.05 | 0.90 – 1.19 |
| Microcephaly, severe (head circumference <3 rd percentile) | 2001-2005 | 599 | 3.18 | 2.93 – 3.44 |
| | 2006-2010 | 846 | 4.23 | 3.95 – 4.52 |
| | 2011-2015 | 1,091 | 5.60 | 5.26 – 5.93 |
| | 2016-2020 | 1,539 | 8.09 | 7.69 – 8.50 |
| Holoprosencephaly | 2001-2005 | 220 | 1.17 | 1.01 – 1.32 |
| | 2006-2010 | 205 | 1.03 | 0.88 – 1.17 |
| | 2011-2015 | 193 | 0.99 | 0.85 – 1.13 |
| | 2016-2020 | 188 | 0.99 | 0.85 – 1.13 |
| Hydrocephaly without spina bifida | 2001-2005 | 1,233 | 6.55 | 6.19 – 6.92 |
| | 2006-2010 | 1,475 | 7.38 | 7.00 – 7.75 |
| | 2011-2015 | 1,678 | 8.61 | 8.19 – 9.02 |
| | 2016-2020 | 1,857 | 9.77 | 9.32 – 10.21 |
| Eye and Ear | | | | |
| Anophthalmia | 2001-2005 | 59 | 0.31 | 0.24 – 0.40 |
| | 2006-2010 | 65 | 0.33 | 0.25 – 0.41 |
| | 2011-2015 | 50 | 0.26 | 0.19 – 0.34 |
| | 2016-2020 | 49 | 0.26 | 0.19 – 0.34 |
| Microphthalmia | 2001-2005 | 502 | 2.67 | 2.43 – 2.90 |
| | 2006-2010 | 590 | 2.95 | 2.71 – 3.19 |
| | 2011-2015 | 515 | 2.64 | 2.41 – 2.87 |
| | 2016-2020 | 525 | 2.76 | 2.52 – 3.00 |
| Cataract | 2001-2005 | 344 | 1.83 | 1.63 – 2.02 |
| | 2006-2010 | 402 | 2.01 | 1.81 – 2.21 |
| | 2011-2015 | 366 | 1.88 | 1.68 – 2.07 |
| | 2016-2020 | 413 | 2.17 | 1.96 – 2.38 |

| Birth Defect (Body System) | 5-Year Period | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|---|---------------|---------------|-------------------|--|
| Anotia or microtia | 2001-2005 | 561 | 2.98 | 2.73 – 3.23 |
| | 2006-2010 | 705 | 3.53 | 3.27 – 3.79 |
| | 2011-2015 | 711 | 3.65 | 3.38 – 3.91 |
| | 2016-2020 | 825 | 4.34 | 4.04 – 4.63 |
| Cardiac and Circulatory | | | | |
| Common truncus | 2001-2005 | 138 | 0.73 | 0.61 – 0.86 |
| | 2006-2010 | 156 | 0.78 | 0.66 – 0.90 |
| | 2011-2015 | 164 | 0.84 | 0.71 – 0.97 |
| | 2016-2020 | 143 | 0.75 | 0.63 – 0.88 |
| Transposition of the great vessels | 2001-2005 | 623 | 3.31 | 3.05 – 3.57 |
| | 2006-2010 | 745 | 3.73 | 3.46 – 3.99 |
| | 2011-2015 | 668 | 3.43 | 3.17 – 3.69 |
| | 2016-2020 | 556 | 2.92 | 2.68 – 3.17 |
| Double outlet right ventricle | 2001-2005 | 386 | 2.05 | 1.85 – 2.26 |
| | 2006-2010 | 450 | 2.25 | 2.04 – 2.46 |
| | 2011-2015 | 484 | 2.48 | 2.26 – 2.70 |
| | 2016-2020 | 505 | 2.66 | 2.42 – 2.89 |
| Tetralogy of Fallot | 2001-2005 | 665 | 3.53 | 3.27 – 3.80 |
| | 2006-2010 | 758 | 3.79 | 3.52 – 4.06 |
| | 2011-2015 | 854 | 4.38 | 4.09 – 4.67 |
| | 2016-2020 | 846 | 4.45 | 4.15 – 4.75 |
| Ventricular septal defect | 2001-2005 | 9,517 | 50.57 | 49.56 – 51.59 |
| | 2006-2010 | 12,698 | 63.51 | 62.41 – 64.62 |
| | 2011-2015 | 12,999 | 66.67 | 65.52 – 67.82 |
| | 2016-2020 | 13,351 | 70.22 | 69.03 – 71.41 |
| Atrial septal defect | 2001-2005 | 9,588 | 50.95 | 49.93 – 51.97 |
| | 2006-2010 | 13,409 | 67.07 | 65.93 – 68.20 |
| | 2011-2015 | 16,816 | 86.25 | 84.94 – 87.55 |
| | 2016-2020 | 17,382 | 91.42 | 90.06 – 92.78 |
| Atrioventricular septal defect (endocardial cushion defect) | 2001-2005 | 762 | 4.05 | 3.76 – 4.34 |
| | 2006-2010 | 887 | 4.44 | 4.14 – 4.73 |
| | 2011-2015 | 891 | 4.57 | 4.27 – 4.87 |
| | 2016-2020 | 851 | 4.48 | 4.17 – 4.78 |
| Pulmonary valve atresia or stenosis | 2001-2005 | 1,530 | 8.13 | 7.72 – 8.54 |
| | 2006-2010 | 2,062 | 10.31 | 9.87 – 10.76 |
| | 2011-2015 | 2,147 | 11.01 | 10.55 – 11.48 |
| | 2016-2020 | 2,206 | 11.60 | 11.12 – 12.09 |
| Tricuspid valve atresia or stenosis | 2001-2005 | 305 | 1.62 | 1.44 – 1.80 |
| | 2006-2010 | 357 | 1.79 | 1.60 – 1.97 |
| | 2011-2015 | 389 | 2.00 | 1.80 – 2.19 |
| | 2016-2020 | 441 | 2.32 | 2.10 – 2.54 |

| Birth Defect (Body System) | 5-Year Period | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|--|---------------|---------------|-------------------|--|
| Ebstein anomaly | 2001-2005 | 137 | 0.73 | 0.61 – 0.85 |
| | 2006-2010 | 131 | 0.66 | 0.54 – 0.77 |
| | 2011-2015 | 155 | 0.79 | 0.67 – 0.92 |
| | 2016-2020 | 165 | 0.87 | 0.74 – 1.00 |
| Aortic valve stenosis | 2001-2005 | 444 | 2.36 | 2.14 – 2.58 |
| | 2006-2010 | 492 | 2.46 | 2.24 – 2.68 |
| | 2011-2015 | 504 | 2.58 | 2.36 – 2.81 |
| | 2016-2020 | 428 | 2.25 | 2.04 – 2.46 |
| Hypoplastic left heart syndrome | 2001-2005 | 391 | 2.08 | 1.87 – 2.28 |
| | 2006-2010 | 435 | 2.18 | 1.97 – 2.38 |
| | 2011-2015 | 485 | 2.49 | 2.27 – 2.71 |
| | 2016-2020 | 439 | 2.31 | 2.09 – 2.52 |
| Patent ductus arteriosus | 2001-2005 | 9,104 | 48.38 | 47.39 – 49.37 |
| | 2006-2010 | 12,187 | 60.95 | 59.87 – 62.04 |
| | 2011-2015 | 14,087 | 72.25 | 71.06 – 73.44 |
| | 2016-2020 | 10,888 | 57.26 | 56.19 – 58.34 |
| Coarctation of the aorta | 2001-2005 | 925 | 4.92 | 4.60 – 5.23 |
| | 2006-2010 | 1,042 | 5.21 | 4.90 – 5.53 |
| | 2011-2015 | 1,094 | 5.61 | 5.28 – 5.94 |
| | 2016-2020 | 1,108 | 5.83 | 5.48 – 6.17 |
| Respiratory | | | | |
| Choanal atresia or stenosis | 2001-2005 | 200 | 1.06 | 0.92 – 1.21 |
| | 2006-2010 | 264 | 1.32 | 1.16 – 1.48 |
| | 2011-2015 | 279 | 1.43 | 1.26 – 1.60 |
| | 2016-2020 | 242 | 1.27 | 1.11 – 1.43 |
| Agenesis, aplasia, or hypoplasia of the lung | 2001-2005 | 623 | 3.31 | 3.05 – 3.57 |
| | 2006-2010 | 537 | 2.69 | 2.46 – 2.91 |
| | 2011-2015 | 531 | 2.72 | 2.49 – 2.96 |
| | 2016-2020 | 482 | 2.53 | 2.31 – 2.76 |
| Oral Clefts | | | | |
| Cleft palate alone (without cleft lip) | 2001-2005 | 1,078 | 5.73 | 5.39 – 6.07 |
| | 2006-2010 | 1,253 | 6.27 | 5.92 – 6.61 |
| | 2011-2015 | 1,110 | 5.69 | 5.36 – 6.03 |
| | 2016-2020 | 1,132 | 5.95 | 5.61 – 6.30 |
| Cleft lip with or without cleft palate | 2001-2005 | 2,062 | 10.96 | 10.48 – 11.43 |
| | 2006-2010 | 2,094 | 10.47 | 10.02 – 10.92 |
| | 2011-2015 | 2,133 | 10.94 | 10.48 – 11.40 |
| | 2016-2020 | 2,013 | 10.59 | 10.12 – 11.05 |
| Gastrointestinal | | | | |
| Tracheoesophageal fistula/ esophageal atresia | 2001-2005 | 381 | 2.02 | 1.82 – 2.23 |
| | 2006-2010 | 412 | 2.06 | 1.86 – 2.26 |
| | 2011-2015 | 460 | 2.36 | 2.14 – 2.57 |
| | 2016-2020 | 443 | 2.33 | 2.11 – 2.55 |

| Birth Defect (Body System) | 5-Year Period | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|---|---------------|---------------|-------------------|--|
| Pyloric stenosis | 2001-2005 | 3,419 | 18.17 | 17.56 – 18.78 |
| | 2006-2010 | 3,818 | 19.10 | 18.49 – 19.70 |
| | 2011-2015 | 2,489 | 12.77 | 12.26 – 13.27 |
| | 2016-2020 | 2,466 | 12.97 | 12.46 – 13.48 |
| Stenosis or atresia of the small intestine | 2001-2005 | 576 | 3.06 | 2.81 – 3.31 |
| | 2006-2010 | 716 | 3.58 | 3.32 – 3.84 |
| | 2011-2015 | 686 | 3.52 | 3.26 – 3.78 |
| | 2016-2020 | 691 | 3.63 | 3.36 – 3.91 |
| Stenosis or atresia of large intestine, rectum, or anal canal | 2001-2005 | 1,009 | 5.36 | 5.03 – 5.69 |
| | 2006-2010 | 1,067 | 5.34 | 5.02 – 5.66 |
| | 2011-2015 | 1,037 | 5.32 | 4.99 – 5.64 |
| | 2016-2020 | 1,026 | 5.40 | 5.07 – 5.73 |
| Hirschsprung disease | 2001-2005 | 239 | 1.27 | 1.11 – 1.43 |
| | 2006-2010 | 284 | 1.42 | 1.26 – 1.59 |
| | 2011-2015 | 307 | 1.57 | 1.40 – 1.75 |
| | 2016-2020 | 297 | 1.56 | 1.38 – 1.74 |
| Biliary atresia | 2001-2005 | 133 | 0.71 | 0.59 – 0.83 |
| | 2006-2010 | 149 | 0.75 | 0.63 – 0.86 |
| | 2011-2015 | 132 | 0.68 | 0.56 – 0.79 |
| | 2016-2020 | 134 | 0.70 | 0.59 – 0.82 |
| Genitourinary | | | | |
| Hypospadias (among males) | 2001-2005 | 5,147 | 53.50 | 52.04 – 54.96 |
| | 2006-2010 | 5,989 | 58.63 | 57.14 – 60.11 |
| | 2011-2015 | 6,714 | 67.35 | 65.74 – 68.96 |
| | 2016-2020 | 6,899 | 71.01 | 69.34 – 72.69 |
| Epispadias | 2001-2005 | 144 | 0.77 | 0.64 – 0.89 |
| | 2006-2010 | 206 | 1.03 | 0.89 – 1.17 |
| | 2011-2015 | 285 | 1.46 | 1.29 – 1.63 |
| | 2016-2020 | 132 | 0.69 | 0.58 – 0.81 |
| Renal agenesis or dysgenesis | 2001-2005 | 997 | 5.30 | 4.97 – 5.63 |
| | 2006-2010 | 1,251 | 6.26 | 5.91 – 6.60 |
| | 2011-2015 | 1,309 | 6.71 | 6.35 – 7.08 |
| | 2016-2020 | 1,534 | 8.07 | 7.66 – 8.47 |
| Bladder exstrophy | 2001-2005 | 41 | 0.22 | 0.16 – 0.30 |
| | 2006-2010 | 35 | 0.18 | 0.12 – 0.24 |
| | 2011-2015 | 36 | 0.18 | 0.13 – 0.26 |
| | 2016-2020 | 21 | 0.11 | 0.07 – 0.17 |
| Musculoskeletal | | | | |
| Congenital hip dislocation without hip dysplasia | 2001-2005 | 581 | 3.09 | 2.84 – 3.34 |
| | 2006-2010 | 537 | 2.69 | 2.46 – 2.91 |
| | 2011-2015 | 389 | 2.00 | 1.80 – 2.19 |
| | 2016-2020 | 297 | 1.56 | 1.38 – 1.74 |

| Birth Defect (Body System) | 5-Year Period | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|--------------------------------------|---------------|---------------|-------------------|--|
| Talipes equinovarus/clubfoot | 2001-2005 | 2,818 | 14.97 | 14.42 – 15.53 |
| | 2006-2010 | 3,117 | 15.59 | 15.04 – 16.14 |
| | 2011-2015 | 3,377 | 17.32 | 16.74 – 17.90 |
| | 2016-2020 | 3,683 | 19.37 | 18.74 – 20.00 |
| Reduction defects of the upper limbs | 2001-2005 | 752 | 4.00 | 3.71 – 4.28 |
| | 2006-2010 | 862 | 4.31 | 4.02 – 4.60 |
| | 2011-2015 | 813 | 4.17 | 3.88 – 4.46 |
| | 2016-2020 | 776 | 4.08 | 3.79 – 4.37 |
| Reduction defects of the lower limbs | 2001-2005 | 364 | 1.93 | 1.74 – 2.13 |
| | 2006-2010 | 403 | 2.02 | 1.82 – 2.21 |
| | 2011-2015 | 391 | 2.01 | 1.81 – 2.20 |
| | 2016-2020 | 364 | 1.91 | 1.72 – 2.11 |
| Craniosynostosis | 2001-2005 | 811 | 4.31 | 4.01 – 4.61 |
| | 2006-2010 | 1,076 | 5.38 | 5.06 – 5.70 |
| | 2011-2015 | 1,232 | 6.32 | 5.97 – 6.67 |
| | 2016-2020 | 1,291 | 6.79 | 6.42 – 7.16 |
| Achondroplasia | 2001-2005 | 53 | 0.28 | 0.21 – 0.37 |
| | 2006-2010 | 69 | 0.35 | 0.27 – 0.44 |
| | 2011-2015 | 72 | 0.37 | 0.29 – 0.47 |
| | 2016-2020 | 91 | 0.48 | 0.39 – 0.59 |
| Diaphragmatic hernia | 2001-2005 | 483 | 2.57 | 2.34 – 2.80 |
| | 2006-2010 | 587 | 2.94 | 2.70 – 3.17 |
| | 2011-2015 | 535 | 2.74 | 2.51 – 2.98 |
| | 2016-2020 | 543 | 2.86 | 2.62 – 3.10 |
| Omphalocele | 2001-2005 | 391 | 2.08 | 1.87 – 2.28 |
| | 2006-2010 | 414 | 2.07 | 1.87 – 2.27 |
| | 2011-2015 | 423 | 2.17 | 1.96 – 2.38 |
| | 2016-2020 | 432 | 2.27 | 2.06 – 2.49 |
| Gastroschisis | 2001-2005 | 843 | 4.48 | 4.18 – 4.78 |
| | 2006-2010 | 1,208 | 6.04 | 5.70 – 6.38 |
| | 2011-2015 | 1,123 | 5.76 | 5.42 – 6.10 |
| | 2016-2020 | 862 | 4.53 | 4.23 – 4.84 |
| Chromosomal | | | | |
| Trisomy 21 (Down syndrome) | 2001-2005 | 2,401 | 12.76 | 12.25 – 13.27 |
| | 2006-2010 | 2,789 | 13.95 | 13.43 – 14.47 |
| | 2011-2015 | 2,760 | 14.16 | 13.63 – 14.68 |
| | 2016-2020 | 2,842 | 14.95 | 14.40 – 15.50 |
| Trisomy 13 (Patau syndrome) | 2001-2005 | 207 | 1.10 | 0.95 – 1.25 |
| | 2006-2010 | 250 | 1.25 | 1.10 – 1.41 |
| | 2011-2015 | 222 | 1.14 | 0.99 – 1.29 |
| | 2016-2020 | 197 | 1.04 | 0.89 – 1.18 |

| Birth Defect (Body System) | 5-Year Period | Cases (count) | Prevalence (rate) | Confidence Interval (95% for Prevalence) |
|--|---------------|---------------|-------------------|--|
| Trisomy 18 (Edwards syndrome) | 2001-2005 | 436 | 2.32 | 2.10 – 2.53 |
| | 2006-2010 | 551 | 2.76 | 2.53 – 2.99 |
| | 2011-2015 | 519 | 2.66 | 2.43 – 2.89 |
| | 2016-2020 | 457 | 2.40 | 2.18 – 2.62 |
| Infants and fetuses with regular reportable birth defects | 2001-2005 | 70,552 | 374.92 | 372.15 – 377.68 |
| | 2006-2010 | 91,187 | 456.08 | 453.12 – 459.04 |
| | 2011-2015 | 107,440 | 551.05 | 547.75 – 554.34 |
| | 2016-2020 | 126,608 | 665.86 | 662.20 – 669.53 |

Prevalence (rate) is expressed as the number of cases per 10,000 live births.

Please see the Methods section of the Annual Report for additional information:

<https://www.dshs.texas.gov/sites/default/files/birthdefects/annualreport/1999-2020-TBDR-Methods.pdf>

Prepared by: Texas Birth Defects Registry, Birth Defects Epidemiology and Surveillance Branch, Texas Department of State Health Services, February 2024.