

2012 HAI Antibiogram, HSR 11

| | aminoglycosides | | | b-lactam/b-lactamase inhibitors | | cephalosporins | | quino-lones | | penicillins | | lincosamide | macrolide | ansamycin | Tetracycline | folate inhibitor | glycopeptide |
|----------------------------|---------------------|----------------------------|--------------------|---------------------------------|-------------------------|---------------------|--------------------|--------------------|---------------------|---------------------|-----------------------|--------------------|--------------------|--------------------|--------------------|-------------------------------|---------------------|
| Pathogen Name | Gentamicin | Gentamicin-High Level Test | Tobramycin | Ampicillin/Sulbactam | Piperacillin/Tazobactam | Cefazolin | Ceftriaxone | Ciprofloxacin | Levofloxacin | Ampicillin | Oxacillin/Methicillin | Clindamycin | Erythromycin | Rifampin | Tetracycline | Trimethoprim/Sulfamethoxazole | Vancomycin |
| Gram-Positive | | | | | | | | | | | | | | | | | |
| Enterococcus faecalis | | (13) 54% | | | | | | | | (14) 100% | | | | | | | (16) 100% |
| Staphylococcus aureus | (26) 88% | | | | | | | (11) 55% | (16) 63% | | (41) 51% | (36) 61% | (38) 34% | (25) 96% | (35) 83% | (32) 94% | (38) 100% |
| Staphylococcus epidermidis | | | | | | | | | | | | | | | | | (26) 96% |
| Gram-Negative | | | | | | | | | | | | | | | | | |
| Klebsiella pneumoniae | (14) 100% | | | (11) 100% | | (11) 100% | (11) 91% | | (11) 100% | (13) 15% | | | | | | (11) 73% | |
| Pseudomonas aeruginosa | | | (11) 82% | | (11) 100% | | | (11) 73% | | | | | | | | | |

Note: Pathogens with less than 10 isolates and antibiotics with less than 25 isolates were excluded from this antibiogram. The number in parentheses indicates the total number of isolates and the percent susceptible is shown in bold).