

# 2012 Diabetes Fact Sheet—Texas Health Service Region 4

## Texas Department of State Health Services

Updated February 2015

### Burden of Disease and Disease Risk Factors

#### Diabetes Prevalence, Adults, 18 years and older

Percentage of Adults (95% confidence interval)

	Region	Texas
Prediabetes	8.3(4.9-13.7)	6.2 (5.3-7.2)
Diabetes	11.4 (8.6-15.0)	10.6 (9.8-11.5)
Overweight and Obese	68.2 (61.7-74.0)	65.1 (63.6-66.5)
No Leisure Time Physical Activity	31.4 (26.1-37.2)	27.2 (25.9-28.6)

An estimated 11.4% of adults in Health Service Region 4 had diabetes and 8.5% had prediabetes.

Prediabetes, diabetes, overweight/obesity and no leisure time physical activity were as common in the region as they were statewide.

#### Age-Adjusted Death Rates, All Ages

Annual Deaths per 100,000 Persons (95% confidence interval)

	Region	Texas
Total	22.4(19.9-24.9)	21.9(21.3-22.5)
White	19.3(16.8-21.9)	16.5(15.8-17.1)
Black	49.7(38.8-60.7)	38.9(36.2-41.5)
Hispanic	--	33.0(31.5-34.6)
Other	--	12.2(9.9-14.4)
Men	23.9(20.1-27.8)	24.0(23.1-24.9)
Women	21.5(18.2-24.8)	20.1(19.3-20.9)

The risk of dying from diabetes was not different in the region than in the state.

The risk of dying from diabetes was not different for whites and blacks in the region than in the state.

The risk of dying from diabetes was not different for men or women in the region than in the state.

#### Age-Adjusted Hospitalization Rates, All Ages

Annual Hospital Discharges per 10,000 Persons (95% confidence interval)

	Region	Texas
Total	16.7(15.9-17.4)	17.2(17.0-17.3)
White	12.4(11.6-13.2)	13.1(12.9-13.3)
Black	32.5(29.8-35.1)	29.2(28.5-29.8)
Hispanic	18.2(15.1-21.4)	18.8(18.5-19.1)
Other	66.3(56.2-76.3)	34.7(33.6-35.9)
Men	17.0(16.0-18.1)	18.5(18.3-18.8)
Women	16.4(15.3-17.4)	16.0(15.8-16.2)
0-17	4.0(3.2-4.7)	3.8(3.6-3.9)
18-44	14.9(13.7-16.1)	12.8(12.6-13.0)
45-64	28.2(26.2-30.1)	27.6(27.2-28.0)
65-74	24.0(21.0-27.0)	37.3(36.4-38.3)
75+	32.1(28.2-36.0)	42.6(41.4-43.8)

For every 10,000 persons, about 17 diabetes hospitalizations occurred in the region and in the state.

Diabetes hospitalizations were higher for the 'Other' race/ethnicity group in the region than in the state.

Diabetes hospitalizations were not different for men or women in the region than in the state.

Diabetes hospitalizations were higher for persons aged 18-44 years in the region than in the state.

Cost Burden


Medicaid Spending among Beneficiaries with Diabetes, All Ages  
For Fee-for-Service and Star & StarPlus Programs

Type of Care	Region			Texas		
	Reimbursement Amount	*Number of Beneficiaries	Average Reimbursement per Beneficiary	Reimbursement Amount	*Number of Beneficiaries	Average Reimbursement per Beneficiary
<b>Total</b>	\$5,591,255.33	12,998	\$430.16	\$279,060,483.30	251,988	\$1,107.44
<b>Inpatient</b>	\$1,997,827.66	401	\$4,982.11	\$40,458,694.90	8,028	\$5,039.70
<b>Outpatient</b>	\$1,044,144.90	16,923	\$175.37	\$16,287,265.74	86,930	\$187.36
<b>Physician</b>	\$2,549,282.77	12,060	\$211.38	\$222,314,522.70	239,783	\$927.15

Medicaid spent \$430.16 per beneficiary with diabetes in the region, an amount less than half the state average. Medicaid spent less on inpatient hospital care, outpatient hospital care and physician care per beneficiary in the region than statewide.

Demography

Total Population = 1,132,414  
Distribution of Population (% of Total Population)



Data Sources: 2012 Texas Vital Statistics, Population Data. Texas Behavioral Risk Factor Surveillance System Public Use Data File, 2013. Texas Department of State Health Services, Center for Health Statistics, Austin, Texas. 2012 Texas Vital Statistics, Mortality Data. 2012 Texas Hospital Inpatient Discharge Public Use Data File. 2012 Texas Medicaid Reimbursement Data as prepared by Research Team, Strategic Decision Support, Texas Health and Human Services Commission, Sep. 2013.

Case Definitions: Prevalence based on respondents 18 years and older who self-report (1) diagnosis of prediabetes, not during pregnancy; (2) diabetes; not during pregnancy (3) body mass index of 25 or greater calculated from height and weight; (4) not participating in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise. Mortality based on ICD-10 E Codes for diabetes (E10-E14). Hospitalizations based on ICD-9 codes for type 1 diabetes (250.01, 250.03, 250.11, 250.13, 250.21, 250.23, 250.31, 250.33, 250.41, 250.43, 250.51, 250.53, 250.61, 250.63, 250.71, 250.73, 250.81, 250.83, 250.91, 250.93), for type 2 diabetes (250.00, 250.02, 250.10, 250.12, 250.20, 250.22, 250.30, 250.32, 250.40, 250.42, 250.50, 250.52, 250.60, 250.62, 250.70, 250.72, 250.80, 250.82, 250.90, 250.92) and diabetes (250.00-250.93). Medicaid reimbursement based on paid and partially paid claims for fee-for-service and primary care case management services selected from the Texas Medicaid and Health Partnership (TMHP) Ad Hoc Query Platform (AHQP) Claims Universe of persons with a primary diagnosis of diabetes (250.00-250.93).

A beneficiary may receive more than one type of care; therefore, the sum of beneficiaries receiving each type of care does not equal the total number of beneficiaries.

Note: "-" indicates too few cases occurred, the sample size was too small, or the relative standard error was >30.0% to provide a reliable estimate.

Statistical significance based upon evaluation of overlap among confidence intervals.