

2010 Texas Childcare Immunization Assessment Coverage Level Results:

This survey is conducted triennially to assess vaccination coverage levels of children attending a childcare facility and to monitor compliance with new and existing vaccine requirements at licensed child care centers (LCCC) and registered child care homes (RCCH) in Texas. A random sample of childcare facilities is selected for the survey every other year. Local health departments and DSHS staff review a random sample of immunization records from each selected childcare facility. Statewide and regional vaccination coverage levels are available for children ages 19 – 59 months of age attending childcare facilities in Texas.

Individual Vaccination Coverage Levels among Childcare Attendees Aged 19 – 59 Months, Texas and HSR, 2010

Area	DTaP‡	Polio§	MMR	Var††	HepB*	Hib¶	PCV§§	HepA**
Texas	89.8	96.0	96.1	95.3	93.5	94.5	86.8	71.3
HSR 1	87.0	94.7	95.4	94.4	93.0	95.1	86.2	66.0
HSR 2/3	83.3	93.0	92.2	91.6	89.4	92.1	81.0	56.2
HSR 4/5N	88.5	95.5	95.4	94.7	94.2	95.0	84.3	72.1
HSR 6/5S	92.2	96.6	97.6	96.5	93.9	94.0	88.8	75.0
HSR 7	91.6	96.5	96.1	95.4	94.2	93.9	88.9	74.7
HSR 8	92.9	97.5	97.8	96.7	96.0	96.7	88.3	76.9
HSR 9/10	91.7	97.8	97.4	96.7	97.4	96.7	89.6	81.9
HSR 11	90.8	97.3	97.2	97.0	94.9	97.4	87.9	76.8

‡ 4 or more doses of any diphtheria and tetanus toxoids and pertussis vaccines including diphtheria and tetanus toxoids, and any acellular pertussis vaccine (DTaP/DTP/DT).

§ 3 or more doses of any poliovirus vaccine.

|| 1 or more doses of measles-mumps-rubella vaccine.

¶ 3 or more doses of Haemophilus influenzae type b (Hib) vaccine.

* 3 or more doses of hepatitis B vaccine.

†† 1 or more doses of varicella at or after child's first birthday, unadjusted for history of varicella illness.

§§ 4 or more doses of PCV.

** 2 or more doses of hepatitis A vaccine.