


# DASHBOARD on Prenatal Oral Health Care

The Pregnancy Risk Assessment Monitoring System (PRAMS) surveys recent mothers about their health practices before, during, and after pregnancy. This information is then used to develop public health programs and policies to ensure that women have healthier pregnancies and healthier babies.

Oral health during pregnancy is critical for ensuring the health of both mother and child.

This DASHBOARD presents information from PRAMS on dental cleanings before and during pregnancy so that the gaps in prenatal oral health care can be identified and targeted.

### Dental Cleaning Before/During Pregnancy by Race/Ethnicity



### Dental Cleaning by Pregnancy Intention



### Dental Cleaning by Payer Source

		Percent	95% Lower CI	95% Upper CI
Before and During Pregnancy	Private Insurance	60.6	57.7	63.4
	Medicaid	38.7	36.0	41.3
	Self pay, Other, Unknown	35.4	30.9	40.0

### Dental Cleaning by Health Service Region


Before and During Pregnancy	HSR	Percent
	HSR 1	35.6
	HSR 2/3	46.5
	HSR 4/5N	41.1
	HSR 6/5S	49.1
	HSR 7	46.6
	HSR 8	50.0
	HSR 9/10	45.4
	HSR 11	40.5

### Dental Cleaning by Income


Before and During Pregnancy	Income Level	Percent
	<15K	35.9
	> 15K to <25K	39.6
	> 25K to <50K	47.2
	> 50K	64.3

# D A S H B O A R D on Prenatal Oral Health Care

**Dental Cleaning by WIC Status**



**Dental Cleaning by Education**



**- 95 Percent Confidence Interval (CI) -**

Each confidence interval presented here is a measure of the precision of its associated prevalence.

Since the prevalence was calculated from a sample of the population, it is an *estimate* of the true value of the population.

A larger sample size will result in a more precise prevalence estimate, and thus, a narrower confidence interval.

If confidence intervals for two estimates do *not* overlap, then there is a statistically significant difference between the prevalence estimates.

However, if confidence intervals *do* overlap, then there may or may not be a statistically significant difference between the prevalence estimates.

**Dental Cleaning by Marital Status**

		Percent	95% Lower CI	95% Upper CI
Before and During Pregnancy	Married	50.3	48.0	52.7
	Not Married	40.3	37.5	43.1

**Dental Cleaning Before/During Pregnancy by Various Characteristics**

	Weighted Frequency	Percent	95% Lower CI	95% Upper CI
Low Birth Weight	36,722	42.7	39.8	45.6
Normal Birth Weight	488,120	46.5	44.5	48.4
Prenatal Care First Trimester	418,954	50.8	48.7	52.9
No Prenatal Care	97,194	33.7	30.2	37.1
Underwt/Normal	295,889	49.6	47.1	52.1
Obese	105,238	40.9	37.1	44.6
Overweight	119,702	44.1	40.4	47.8
Physical Abuse	19,768	31.6	24.5	38.7
No Physical Abuse	474,149	46.6	44.7	48.5

**Dental Cleaning by Mother's Place of Birth**

Before and During Pregnancy	Foreign Born	38.5
	US Born	49.6

