



OB _____
CDC Use Only

REPORT OF OUTBREAK OF SUSPECTED VIRAL GASTROENTERITIS

Viral Gastroenteritis Section	Telephone	(404) 639-3577 or
DASH Unit 75		(404) 639-3607
Centers for Disease Control and Prevention	Facsimile	(404) 639-3645
1600 Clifton Road, N.E., Mailstop G-04	E-Mail	CaliciNet@cdc.gov
Atlanta, GA 30333		

Primary contact for epidemiologic investigation Date _____
mm/dd/yyyy

Name _____ Telephone _____

Agency _____ Facsimile _____

Address _____ E-mail _____

Outbreak Information

State Outbreak ID _____ EFORS code _____ Date health department notified _____
(if known) mm/dd/yyyy

Date of first case _____ Date of last case _____
mm/dd/yyyy mm/dd/yyyy

Location(s) of outbreak: State _____ City _____ County (list if several) _____

If multistate, list other states _____

Suspected mode of transmission (can check more than one)

- Person-to-person Foodborne Waterborne
- Unknown Other _____

If food or waterborne

Implicated food or water source _____ Foodhandler implicated? _____
(can check more than one)

Yes: epidemiologic evidence
 Yes: laboratory evidence
 Yes: suspected, but no evidence
 No

Setting (if there is an additional setting, please add below in comments)

- Nursing home Assisted Living Restaurant / Deli / Cafeteria Hotel School Daycare Camp
- Community Prison Ship Religious Facility Hospital Private event Catered Event
- Work Place Private Home Other _____ Date of event (if applicable) _____
mm/dd/yyyy

Illness Characteristics

Number of persons exposed _____ Number of persons ill _____

Number visiting health care provider _____ Number hospitalized _____ Number deaths _____
(Categories NOT mutually exclusive)

Symptoms: Number of persons with information _____ Median incubation period (hours) _____ range _____

No. with abdominal cramps _____ No. with fever _____

No. with diarrhea _____ No. with vomiting _____ Median duration of illness (hours) _____ range _____

No. with other symptom(s) _____

Comments: _____

RECOMMENDATIONS REGARDING SPECIMEN COLLECTION FOR DIAGNOSIS OF NLVs*

Clinical Specimens

Stool

Timing. Specimen collection for viral testing should begin on day 1 of the epidemiologic investigation. Any delays to await testing results for bacterial or parasitic agents could preclude establishing a viral diagnosis. Ideally, specimens should be obtained during the acute phase of illness (i.e., within 48--72 hours after onset) while the stools are still liquid or semisolid because the level of viral excretion is greatest then. With the development of sensitive molecular assays, the ability to detect viruses in specimens collected later in the illness has been improved. In specific cases, specimens might be collected later during the illness (i.e., 7--10 days after onset), if the testing is necessary for either determining the etiology of the outbreak or for epidemiologic purposes (e.g., a specimen obtained from an ill foodhandler who might be the source of infection). If specimens are collected late in the illness, the utility of viral diagnosis and interpretation of the results should be discussed with laboratory personnel before tests are conducted.

Number and Quantity. Ideally, specimens from ≥ 10 ill persons should be obtained during the acute phase of illness. Bulk samples (i.e., 10--50 ml of stool placed in a stool cup or urine container) are preferred, as are acute diarrhea specimens that are loose enough to assume the shape of their containers. Serial specimens from persons with acute, frequent, high-volume diarrhea are useful as reference material for the development of assays. The smaller the specimen and the more formed the stool, the lower the diagnostic yield. Rectal swabs are of limited or no value because they contain insufficient quantity of nucleic acid for amplification.

Storage and Transport. Because freezing can destroy the characteristic viral morphology that permits a diagnosis by EM, specimens should be kept refrigerated at 4 C. At this temperature, specimens can be stored without compromising diagnostic yield for 2--3 weeks, during which time testing for other pathogens can be completed. If the specimens have to be transported to a laboratory for testing, they should be bagged and sealed and kept on ice or frozen refrigerant packs in an insulated, waterproof container. If facilities for testing specimens within 2--3 weeks are not available, specimens can be frozen for antigen or PCR testing.

Vomit

Vomiting is the predominant symptom among children, and specimens of vomitus can be collected to supplement the diagnostic yield from stool specimens during an investigation. Recommendations for collection, storage, and shipment of vomitus specimens are the same as those for stool specimens.

Serum

Timing. If feasible, acute- and convalescent-phase serum specimens should be obtained to test for a diagnostic ≥ 4 -fold rise in IgG titer to NLVs. Acute-phase specimens should be obtained during the first 5 days of symptoms, and the convalescent-phase specimen should be collected from the third to sixth week after resolution of symptoms.

Number and Quantity. Ideally, 10 pairs of specimens from ill persons (i.e., the same persons submitting stool specimens) and 10 pairs from well persons (controls) should be obtained. Adults should provide 5--7 ml of blood, and children should provide 3--4 ml.

Storage. Specimens should be collected in tubes containing no anticoagulant, and the sera should be spun off and frozen. If a centrifuge is not available, a clot should be allowed to form, and the serum should be decanted and frozen. If this step cannot be accomplished, the whole blood should be refrigerated but not frozen.

Environmental Specimens

NLVs cannot be detected routinely in water, food, or environmental specimens. Nevertheless, during recent outbreaks (33--36), NLVs have been detected successfully in vehicles epidemiologically implicated as the source of infection. If a food or water item is strongly suspected as the source of an outbreak, then a sample should be obtained as early as possible and stored at 4 C. If the epidemiologic investigation confirms the link, a laboratory with the capacity to test these specimens should be contacted for further testing. If drinking water is suspected, special filtration (45) of large volumes (i.e., 5--100 liters) of water can concentrate virus to facilitate its detection.