

TABLE V
REPORTED DISEASES BY AGE GROUP¹
CASES PER 100,000 POPULATION² 2011

DISEASE	<1	1-4	5-9	10-14	15-19	20-29	30-39	40-49	50-59	60+	TOTAL
AMEBIASIS	0.0	0.2	0.2	0.2	0.5	0.6	0.6	0.5	0.5	0.3	0.4
ANAPLASMA PHAGOCYTOPHILUM	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
BOTULISM, INFANT ³	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0
BOTULISM, WOUND	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
BRUCELLOSIS	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.1	0.0
CAMPYLOBACTERIOSIS	23.5	18.5	8.2	5.6	4.8	5.1	4.6	4.9	6.3	6.6	6.7
CHICKENPOX (VARICELLA)	50.6	40.3	37.9	26.5	6.2	3.8	3.0	1.8	1.4	1.3	9.9
CHOLERA	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
CREUTZFELEDT-JAKOB DISEASE	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.3	0.1
CRYPTOSPORIDIOSIS	2.0	5.0	3.1	1.7	1.8	1.7	1.8	1.4	1.2	1.9	1.9
CYCLOSPORIASIS	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.0	0.1	0.1
CYSTICERCOSIS	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.0
DENGUE	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0
EHRLICHIOSIS/ANAPLASMOSIS	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
ENCEPHALITIS, NONARBOVIRAL	0.0	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.2	0.1
ESCHERICHIA COLI, SHIGA TOXIN-PRODUCING (STEC)	12.9	12.8	2.0	1.4	1.7	0.9	0.5	0.6	0.6	1.1	1.9
HAEMOPHILUS INFLUENZAE TYPE B, INVASIVE	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
HEMOLYTIC UREMIC SYNDROME	0.0	0.9	0.2	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.1
HEPATITIS A, ACUTE	0.0	0.1	0.1	0.4	0.6	0.7	0.6	0.4	0.5	1.0	0.5
HEPATITIS B, ACUTE	0.0	0.0	0.1	0.1	0.2	0.7	1.4	1.6	1.1	0.6	0.8
HEPATITIS B, PERINATAL ⁴	0.5	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5
HEPATITIS C, ACUTE	0.2	0.1	0.0	0.0	0.1	0.2	0.1	0.2	0.3	0.1	0.1
HEPATITIS E, ACUTE	0.0	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.0	0.0	0.1
INFLUENZA-ASSOCIATED PEDIATRIC MORTALITY ⁵	1.0	0.2	0.0	0.1	0.2	0.0	0.0	0.0	0.0	0.0	0.2
JAPANESE ENCEPHALITIS	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
LEGIONELLOSIS	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.5	0.8	1.6	0.4
LEISHMANIASIS	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
LISTERIOSIS	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.3	0.8	0.2
LYME DISEASE	0.0	0.0	0.2	0.1	0.3	0.4	0.3	0.3	0.3	0.3	0.3
MALARIA	0.0	0.1	0.2	0.2	0.2	0.7	0.4	0.7	0.4	0.2	0.4
MEASLES	0.5	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
MENINGITIS, ASEPTIC	73.8	2.2	5.0	4.2	4.3	5.5	4.5	3.2	2.7	3.2	5.0
BACTERIAL MENINGITIS	16.4	0.9	0.8	0.6	0.4	0.6	0.7	1.4	1.7	1.9	1.3
OTHER MENINGITIS	0.0	0.1	0.0	0.0	0.2	0.1	0.5	0.7	0.4	0.3	0.3
MENINGOCOCCAL INFECTION	1.0	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.0	0.1	0.1
MUMPS	0.0	0.6	0.5	0.4	0.1	0.5	0.3	0.2	0.1	0.1	0.3
PERTUSSIS	66.7	12.3	8.8	5.0	2.0	0.9	1.6	1.1	1.2	0.8	3.7
Q FEVER, ACUTE	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.2	0.1
Q FEVER, CHRONIC	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
ROCKY MOUNTAIN SPOTTED FEVER	0.0	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.4	0.2	0.2
SALMONELLOSIS	212.8	64.6	26.4	16.1	10.2	10.0	7.7	10.9	13.3	22.1	20.2
SHIGELLOSIS	10.5	57.2	45.7	10.3	3.2	4.0	3.5	2.1	1.7	1.4	9.8
STREPTOCOCCUS, GROUP A	2.2	0.9	0.5	0.6	0.6	0.7	0.9	1.7	2.2	4.8	1.6
STREPTOCOCCUS, GROUP B	31.8	0.0	0.1	0.1	0.1	0.7	1.0	3.0	6.3	10.4	3.5
STREPTOCOCCUS PNEUMONIAE	12.7	6.4	2.4	1.3	0.8	2.0	2.9	5.3	10.3	17.4	6.2
TAENIASIS	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
TETANUS	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
TRICHINOSIS	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
TYPHOID FEVER	0.0	0.1	0.2	0.2	0.2	0.1	0.1	0.0	0.1	0.1	0.1
TYPHUS, MURINE	0.0	0.6	1.1	2.3	1.4	0.8	0.7	1.0	1.4	1.3	1.1
VIBRIO PARAHAEMOLYTICUS	0.0	0.0	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.1	0.1
VIBRIO VULNIFICUS	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.1
VIBRIO, OTHER/UNSPECIFIED	0.0	0.2	0.1	0.3	0.2	0.1	0.1	0.1	0.1	0.2	0.1
VISA	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0
WEST NILE FEVER	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0
WEST NILE NEUROINVASIVE DISEASE	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.3	0.1
YERSINIOSIS	1.5	0.1	0.0	0.0	0.1	0.1	0.1	0.0	0.0	0.1	0.1

¹ Age is calculated based on date of birth and event date, which is onset date if known.

² DSHS Center for Health Statistics projected 2011 Texas population data http://www.dshs.texas.gov/chs/popdat/st2011_p.shtm

³ Infant botulism rates are calculated using the population under 1 year of age.

⁴ Perinatal hepatitis B cases are defined as infants >1 month of age through 24 months of age who were born in the US to HBsAg positive mothers. The rates were calculated using the population under 2 years of age, which approximates this cohort.

⁵ Influenza-associated pediatric mortality is calculated using the population under 18 years of age.